

PRODUCT MONOGRAPH

Pr ACT FAMCICLOVIR

Famciclovir

125 mg, 250 mg and 500 mg Tablets

ANTIVIRAL AGENT

Actavis Pharma Company
6733 Mississauga Road, Suite 400
Mississauga, Ontario
L5N 6J5

Date of Revision:
December 06, 2016

Submission Control No.: 200282

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	5
DRUG INTERACTIONS	9
DOSAGE AND ADMINISTRATION	10
OVERDOSAGE	12
ACTION AND CLINICAL PHARMACOLOGY	12
STORAGE AND STABILITY.....	15
DOSAGE FORMS, COMPOSITION AND PACKAGING	15
PART II: SCIENTIFIC INFORMATION	16
PHARMACEUTICAL INFORMATION.....	16
CLINICAL TRIALS	17
TOXICOLOGY	22
REFERENCES	25
PART III: CONSUMER INFORMATION.....	27

ACT FAMCICLOVIR

(famciclovir)

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All non-medicinal ingredients
Oral	Tablets/ 125 mg, 250 mg and 500 mg	hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium starch glycolate, talc, and titanium dioxide

INDICATIONS AND CLINICAL USE

ACT FAMCICLOVIR (famciclovir) tablets are indicated:

- For the treatment of acute herpes zoster (shingles);
- For the treatment or suppression of recurrent episodes of genital herpes in immunocompetent adults;
- For the treatment of recurrent episodes of mucocutaneous herpes simplex infections in HIV-infected patients.

Early treatment of acute herpes zoster (shingles) in immune-competent individuals with oral famciclovir resulted in decreased time to loss of vesicles; decreased time to loss of crusts; and decreased viral shedding.

The results of clinical studies indicate that early treatment of acute herpes zoster with oral famciclovir resulted in decreased duration of post-herpetic neuralgia. Those most likely to benefit are patients who initiate treatment within 48 hours of onset of rash or are greater than 50 years of age or those patients with severe pain at the time of treatment initiation.

In clinical studies of immunocompetent patients with recurrent genital herpes (typically ≥ 6 episodes in a 12 month period) famciclovir suppressed lesional episodes, slowed the rate to first recurrence and patients were more likely to remain free from recurrences for a 12-month period. Suppressive therapy in patients with fewer than 6 episodes of genital herpes in a 12 month period was not evaluated in these clinical studies.

Initiation of famciclovir treatment of recurrent genital herpes during the prodrome or as soon as possible after the onset of lesions resulted in decreased duration of viral shedding, decreased time to lesion healing and decreased time to resolution of symptoms (including pain, tenderness, itching and burning).

CONTRAINDICATIONS

Patients who have known hypersensitivity to famciclovir or to any ingredient in the formulation or component of the container. For a complete listing, see the Dosage Forms, Composition and Packaging section of this product monograph.

WARNINGS AND PRECAUTIONS

General

The efficacy of famciclovir has not been established for first episode genital herpes infections, disseminated zoster, or in immunocompromised patients with herpes zoster (see ACTION AND CLINICAL PHARMACOLOGY). Dosage adjustment is required when administering famciclovir to patients with moderate or severe renal dysfunction (see DOSAGE AND ADMINISTRATION). No special precautions are required for patients with mild or moderate hepatic impairment. Famciclovir has not been studied in patients with severe hepatic impairment (see ACTION AND CLINICAL PHARMACOLOGY). Conversion of famciclovir to the active metabolite penciclovir may be impaired in these patients resulting in lower penciclovir plasma concentrations, and thus possibly a decrease of efficacy of famciclovir (see ACTION AND CLINICAL PHARMACOLOGY).

Genital herpes is a sexually transmitted disease with an increased risk of transmission during acute episodes. There are no data evaluating whether famciclovir will prevent transmission of infection to others. Patients should be advised to avoid intercourse when lesions and/or symptoms are present (even if treatment with an anti-viral has been initiated) in order to avoid infecting partners. Genital herpes can also be transmitted in the absence of symptoms through asymptomatic viral shedding.

Pregnancy:

Although animal studies have not shown any embryotoxic or teratogenic effects with famciclovir or penciclovir, the active metabolite of famciclovir, there are insufficient data on the use of famciclovir in pregnant women. Because animal reproductive studies are not always predictive of human response, ACT FAMCICLOVIR should, therefore, not be used in pregnancy unless the potential benefits are considered to outweigh the potential risks associated with treatment.

Breast-feeding:

Following oral administration of famciclovir to lactating rats, penciclovir was excreted in breast milk. It is not known whether it (penciclovir) is excreted in human milk, thus, a decision should

be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Use in Children:

Safety and efficacy in children under the age of 18 years have not been established.

Race:

The impact of race on the safety and efficacy of famciclovir has not been established.

Geriatric Use:

Of 816 patients with herpes zoster in clinical studies who were treated with famciclovir, 248 (30.4%) were > 65 years of age and 103 (13%) were > 75 years of age. No overall differences were observed in safety between younger and older patients (see Adverse Events).

Impairment of Fertility

As with other drugs of this class, testicular toxicity has been observed in animals receiving both famciclovir and penciclovir. Famciclovir has been shown to have no significant effects on sperm count, morphology, or motility in man. Clinical data do not indicate an impact of famciclovir on male fertility following long-term treatment at an oral dose of 250 mg twice daily.

Effects on ability to drive and use machines:

ACT FAMCICLOVIR can cause dizziness, drowsiness or confusion in very rare cases. Patients who experience any of these symptoms while taking ACT FAMCICLOVIR should take special care when driving or using machines (see ADVERSE REACTIONS – Post-Market Adverse Drug Reactions).

ADVERSE REACTIONS

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adverse Drug Reaction Overview

Immunocompetent Patients

The most frequent adverse reactions reported during herpes zoster clinical trials with oral famciclovir three times daily were as shown in the following table.

Patients (%) reporting A/Es related* to study medication by preferred term in Famciclovir Zoster trials within 30 days of the last dose.

	Famciclovir	Placebo
Patients receiving study medication	816	146
Event:	%	%
Body as a Whole		
Headache	7.1	6.8
Fatigue	1.6	0.7
Fever	1.1	0.0
Rigors	0.6	1.4
Herpes Zoster Symptoms	0.5	1.4
Central Nervous System		
Dizziness	1.5	0.7
Somnolence	1.2	2.7
Gastrointestinal		
Nausea	4.3	8.2
Diarrhea	1.8	2.1
Abdominal Pain	1.5	0.0
Constipation	1.0	0.0
Vomiting	1.2	0.7
Anorexia	0.5	1.4
Dermatologic		
Pruritis	1.2	0.7
Sweating increased	1.0	0.0
Hepatic		
ALT (SGPT) Increased	0.6	1.4
Gamma GT Increased	0.6	1.4
Hepatic Enzymes Increased	0.2	1.4
Special Senses		
Tinnitus	0.0	1.4

* Includes events assessed by the investigator as related, probably related, possibly related and AEs where the relationship was unassessable or missing.

The most frequent adverse reactions reported within 30 days of the last dose, during genital herpes clinical trials with oral famciclovir were as shown in the following table.

Patients (%) reporting A/Es related* to study medication by preferred term in Famciclovir Genital Herpes trials.

	Famciclovir	Placebo
Patients receiving study medication	1500	255
Event:	%	%
Body as a Whole		
Headache	5.5	3.9
Fatigue	1.5	1.6
Central Nervous System		
Dizziness	2.3	3.1
Gastrointestinal		
Nausea	4.9	3.9
Diarrhea	1.8	1.6
Dyspepsia	1.3	1.2
Abdominal Pain	0.9	1.6
Autonomic Nervous System		
Mouth Dry	0.3	1.2

* Includes events assessed by the investigator as related, probably related, possibly related or where relationship was unassessable or not given.

The most frequent adverse events (incidence of >1%) are listed in the following table for patients receiving double-blind famciclovir or placebo for at least 10 months in the two 12-month-long trials.

Patients (%) reporting A/Es related* to study medication by preferred term in Famciclovir Genital Herpes Suppression trials.

	Famciclovir	Placebo
Patients receiving study medication	458	63
Event:	%	%
Body as a Whole		

	Famciclovir	Placebo
Patients receiving study medication	458	63
Headache	8.7	9.5
Central Nervous System		
Dizziness	1.5	0
Gastrointestinal		
Abdominal Pain	2.4	4.8
Dyspepsia	2.0	3.2
Nausea	1.5	3.2
Diarrhea	1.3	0
Flatulence	1.1	0
Enzyme Abnormality [†]	2.2	3.2
Bilirubinemia	1.3	1.6
Leukopenia	1.3	0

* Includes events assessed by the investigator as related, probably related, possibly related and AEs where the relationship was unassessable or missing.

[†] Reports of elevated lipase.

HIV-Infected Patients

In a controlled study of HIV-infected patients, the overall percentages of patients reporting adverse events were comparable for famciclovir and acyclovir. The most frequently reported events ($\geq 2\%$ in any group) are listed below.

Adverse events* reported by preferred term in HIV-infected patients

	Famciclovir	Acyclovir
Patients receiving study medication	150	143
Event:	%	%
Headache	13.3	9.1
Nausea	8.7	8.4
Diarrhea	4.7	4.9
Vomiting	3.3	2.1

	Famciclovir	Acyclovir
Patients receiving study medication	150	143
Fatigue	2.0	0.7
Creatine phosphokinase increased	2.0	0.7
Abdominal Pain	1.3	3.5

* Includes adverse events considered by the investigators to be related, possibly related or of unknown relationship to study medication.

Post-Market Adverse Drug Reactions

Because they are reported spontaneously from a population of unknown size, estimates of frequency of post-marketing adverse events cannot be made. The following events have been chosen for inclusion due to their seriousness, frequency of reporting, potential causal connection to famciclovir, or a combination of these factors headache, abdominal pain, diarrhea, nausea and confusion (including delirium, disorientation, confusional state, occurring predominantly in the elderly), rash, urticaria, pruritus, serious skin reactions (e.g. erythema multiforme, Steven Johnson syndrome, toxic epidermal necrolysis, leukocytoclastic vasculitis), angioedema (e.g. face edema, eyelid edema, periorbital edema, pharyngeal edema), vomiting, dizziness, somnolence (predominantly in the elderly), hallucinations, palpitations, jaundice and abnormal liver function tests, seizure, anaphylactic shock and anaphylactic reaction.

Abnormal Hematological and Clinical Chemistry Findings: In post-market experience, thrombocytopenia has been reported.

DRUG INTERACTIONS

Effects of other medicinal products on famciclovir

No clinically significant alterations in penciclovir (active metabolite of famciclovir) pharmacokinetics were observed following single dose administration of 500 mg famciclovir after pretreatment with multiple doses of cimetidine, allopurinol, theophylline, zidovudine, or promethazine or when given shortly after an antacid (magnesium and aluminium hydroxide) or concomitantly with emtricitabine. Furthermore, no clinically significant effect on penciclovir pharmacokinetics was observed following multiple-dose (three times daily) administration of famciclovir (500 mg) with multiple doses of digoxin.

Concurrent use of probenecid may result in increased plasma concentrations of penciclovir the active metabolite of famciclovir (see ACTION and CLINICAL PHARMACOLOGY). Therefore, patients receiving ACT FAMCICLOVIR at a dose of 500 mg three times daily co-

administered with probenecid on consecutive days should be monitored for toxicity and dose reduction of ACT FAMCICLOVIR may be considered.

The conversion of the inactive metabolite 6-deoxy penciclovir (formed by deacetylation of famciclovir) to penciclovir is catalysed by aldehyde oxidase. Interactions with other drugs metabolized by this enzyme and/or inhibiting this enzyme could potentially occur. Clinical interaction studies of famciclovir with cimetidine and promethazine, in *vitro* inhibitors of aldehyde oxidase, did not show relevant effects on the formation of penciclovir. However raloxifene, the most potent aldehyde oxidase inhibitor observed in *vitro*, could affect the formation of penciclovir and thus the efficacy of famciclovir. When raloxifene is co-administered with famciclovir the clinical efficacy of the antiviral therapy should be monitored.

Effects of famciclovir on other medicinal products

The pharmacokinetics of digoxin were not altered by concomitant administration of single or multiple (three times daily) doses of famciclovir (500 mg). No clinically significant effects on the pharmacokinetics of zidovudine, its metabolite zidovudine glucuronide or emtricitabine were observed following a single oral dose of 500 mg famciclovir co-administered with zidovudine or emtricitabine.

Although famciclovir is only a weak inhibitor of aldehyde oxidase in *vitro*, interactions with drugs metabolized by aldehyde oxidase could potentially occur. Evidence from preclinical studies has shown no potential for induction of cytochrome P450 enzymes and inhibition of CYP3A4.

DOSAGE AND ADMINISTRATION

Recommended Dose:

Herpes zoster infections:

The recommended dose is 500 mg 3 times per day for 7 days. Therapy should be initiated within 72 hours of the onset of the rash.

Herpes simplex infections:

Immunocompetent Patients

Recurrent genital herpes episodes: The recommended dosage is 125 mg twice a day for 5 days. Initiation of treatment is recommended during the prodromal period or as soon as possible at the first sign or symptom of a recurrent episode (e.g. tingling, itching, burning, pain, or lesion).

Suppression of recurrent genital herpes episodes: The recommended dosage is 250 mg twice daily for up to 1 year. The safety and efficacy of famciclovir therapy beyond one year of treatment has not been established.

HIV-Infected Patients

For recurrent episodes of mucocutaneous herpes simplex infection, the recommended dosage is 500 mg twice a day for 7 days. Treatment should be initiated at the first sign or symptom of a recurrent episode (e.g. tingling, itching, burning, pain, or lesion).

ACT FAMCICLOVIR tablets should be swallowed whole and may be taken with or without food.

Dosage Adjustment

Dosage in renally impaired patients:

In patients with moderately or severely reduced renal function, dosage reduction is recommended:

<u>Indication</u>	<u>Creatinine clearance (mL/min/1.73m²)</u>	<u>Dosage</u>
Herpes Zoster	>60	500 mg every 8 hours
	40-59	500 mg every 12 hours
	20-39	500 mg every 24 hours
	<20	250 mg every 24 hours
	hemodialysis patients	250 mg following each dialysis during 7 days
Recurrent Genital Herpes	>20	125 mg every 12 hours
	<20	125 mg every 24 hours
	hemodialysis patients	125 mg following each dialysis during 5 days
Suppression of Recurrent Genital Herpes	>40	250 mg every 12 hours
	20-39	125 mg every 12 hours
	<20	125 mg every 24 hours
	hemodialysis patients	125 mg following each dialysis
Recurrent episodes of mucocutaneous herpes simplex infections in HIV-infected patients	>40	500 mg every 12 hours
	20-39	500 mg every 24 hours
	<20	250 mg every 24 hours
	hemodialysis patients	250 mg following each dialysis during 7 days

Dosage in hepatically impaired patients:

No dose adjustment is required in patients with mild or moderate hepatic impairment. No data are available for patients with severe hepatic impairment (see ACTION and CLINICAL PHARMACOLOGY).

Missed Dose

If a dose of ACT FAMCICLOVIR is missed, it should be taken as soon as the patient remembers. The next dose should be taken at the normal time.

The patient should carry on as normal until they have finished all the tablets.

Do not double-dose.

OVERDOSAGE

For management of suspected overdose, contact the regional poison control centre.

No acute overdosage has been reported. Appropriate symptomatic and supportive therapy should be given. Penciclovir, the active metabolite of famciclovir, is dialyzable; plasma concentrations are reduced by approximately 75% following 4 h hemodialysis.

In patients with underlying renal disease who have received inappropriately high doses of famciclovir for their level of renal function, acute renal failure has been reported frequently.

ACTION AND CLINICAL PHARMACOLOGY

Famciclovir is the orally administered pro-drug of the antiviral agent penciclovir. Famciclovir itself has no antiviral activity until it is biotransformed to penciclovir. Studies in volunteers have shown that famciclovir is well absorbed and produces plasma penciclovir concentrations superior to those obtained following oral administration of penciclovir alone.

The mean bioavailability of penciclovir after administration of oral famciclovir is 77%. The mean peak plasma concentration of penciclovir, following a 500 mg oral dose of famciclovir was 3.3 mcg/mL and occurred at a mean time of 0.89 hours post-dose. Plasma concentration time curves of penciclovir are similar following single and repeat dosing. The terminal plasma elimination half-life of penciclovir after both single and repeat oral dosing with famciclovir is 2.3 hours. The elimination of famciclovir is by metabolism, principally to penciclovir and its 6-deoxy precursor, which are subsequently excreted in urine (see Pharmacokinetics).

Mechanism of Action

Penciclovir is a substituted guanine analogue with potent and selective antiviral activity against varicella zoster virus and other human herpes viruses (see Virology). Penciclovir is in the same class of antiviral drugs as acyclovir, and both are phosphorylated by viral thymidine kinase and then by cellular kinases to the active triphosphate form in virus-infected cells. Penciclovir triphosphate inhibits viral DNA polymerase competitively with deoxyguanosine triphosphate and is incorporated into the extending DNA chain, preventing significant chain elongation. Consequently, viral DNA synthesis and, therefore, viral replication are inhibited. Inhibition of the virus reduces the period of viral shedding, limits the degree of spread and level of pathology, and thereby facilitates healing.

Penciclovir is not readily phosphorylated in uninfected cells and does not inhibit cellular DNA synthesis even at concentrations > 20 times those achieved in clinical usage.

Pharmacokinetics

Absorption: Following oral administration, famciclovir is rapidly, extensively and consistently absorbed and converted to the antivirally active compound, penciclovir. The mean (range) bioavailability of penciclovir after oral famciclovir is 77% (69.5 – 84.5%). The extent of systemic availability (AUC) of penciclovir from oral famciclovir is unaffected by food. In a healthy male volunteer study using a single oral dose of famciclovir, the pharmacokinetics of penciclovir were linear over the famciclovir dose range 125 to 750 mg. The mean (range) peak plasma concentration of penciclovir, calculated from dose normalised estimates across all single dose healthy male volunteer studies, following a single 500 mg dose of famciclovir was 3.3 ug/ml (range 1.3 – 6.3 ug/ml) and occurred at a mean time of 0.89 hours post-dose (range 0.5 – 5.0 hours). The mean terminal half-life of penciclovir was 2.3 hours (range 0.99 – 5.26 hours).

Pharmacokinetic parameter estimates of penciclovir following oral administration of a single dose of famciclovir to patients with uncomplicated herpes zoster were essentially identical to values reported in healthy volunteers matched for age. Repeated oral dosing of famciclovir every 8 hours for up to 7 days in patients with herpes zoster infections had no significant effect on the pharmacokinetics of penciclovir compared to that described after single doses of famciclovir. The terminal plasma half-life of penciclovir in patients with herpes zoster was 2.8 h and 2.7 h, respectively, after single and repeated doses of famciclovir.

Distribution: Plasma protein binding of penciclovir and its 6-deoxy precursor is low (<20%) and penciclovir distributes freely between plasma and blood cells.

Metabolism: Following oral administration little or no famciclovir is detected in plasma or urine since famciclovir is rapidly converted via deacetylation and oxidation to penciclovir. An *in vitro* study using human liver microsomes demonstrated that cytochrome P450 does not play an important role in famciclovir metabolism. The conversion of B-deoxy penciclovir is catalyzed by aldehyde oxidase.

Excretion: Little or no famciclovir is detected in plasma or urine since famciclovir undergoes extensive first-pass metabolism to penciclovir. The major metabolites identified in plasma and urine are penciclovir ($67 \pm 4\%$ of radioactivity in plasma at 1.5 h following a 500 mg oral dose of [^{14}C] famciclovir and $82 \pm 2.2\%$ of radioactivity in 0-24 h urine) and, to a lesser extent, its 6-deoxy precursor, which has no antiviral activity ($11 \pm 4\%$ in plasma and $7 \pm 0.5\%$ in urine at the corresponding time points). Other minor, virally inactive metabolites identified in human urine are monoacetylated penciclovir and 6-deoxy monoacetylated penciclovir (each $< 0.5\%$ of the dose).

Renal clearance values for penciclovir exceed creatinine clearance indicating that net active tubular secretion and glomerular filtration contribute to renal elimination.

Renal Impaired Patients: The apparent plasma clearance, renal clearance, and plasma elimination rate constant of penciclovir decreased linearly with reductions in renal function, both after single and repeated dosing. Dose adjustment is necessary in patients with renal insufficiency (see DOSAGE AND ADMINISTRATION).

Hepatically Impaired Patients: Following single oral administration of famciclovir to patients with mild or moderate hepatic impairment, there was no change in the extent of availability of penciclovir compared with healthy volunteers. There was, however, a decrease in the rate of availability of penciclovir in the hepatically impaired subjects. Mean maximum plasma concentrations of penciclovir were decreased by 43% and the time to maximum plasma concentrations increased by 0.75 hours. However, no dosage adjustment for patients with mild or moderate hepatic impairment is recommended. The pharmacokinetics of penciclovir following oral famciclovir in patients with severe hepatic impairment has not been studied. Conversion of famciclovir to the active metabolite penciclovir may be impaired in these patients resulting in lower penciclovir plasma concentrations, and thus possibly a decrease of efficacy of famciclovir (see WARNINGS AND PRECAUTIONS).

HIV-Infected Patients: Following oral administration of a single dose of 500 mg famciclovir to HIV-positive patients, the pharmacokinetic parameters of penciclovir were comparable to those observed in healthy subjects.

Elderly Subjects:

Based on cross-study comparisons, the mean penciclovir AUC was about 40 % higher and penciclovir renal clearance about 20% lower after oral administration of famciclovir in elderly volunteers (65-79 years) compared to younger volunteers. Some of this difference may be due to differences in renal function between the two age groups. No dose adjustment based on age is recommended unless renal function is impaired (see Dosage and Administration).

Gender:

Small differences in renal clearance of penciclovir between females and males have been reported and were attributed to gender differences in renal function. No dose adjustment based on gender is recommended.

STORAGE AND STABILITY

ACT FAMCICLOVIR should be stored at room temperature (15-30°C). Protect from light.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Dosage Forms

ACT FAMCICLOVIR is available as 125 mg, 250 mg or 500 mg tablets with the following descriptions:

ACT FAMCICLOVIR 125 mg tablets are white to off-white, film-coated, round biconvex tablet with “FC” on one side and “>” on the other.

125

ACT FAMCICLOVIR 250 mg tablets are white to off-white, film-coated, round biconvex tablet with “FC” on one side and “>” on the other.

250

ACT FAMCICLOVIR 500 mg tablets are white to off-white, film-coated, capsule-shaped, biconvex tablet with “FC500” on one side and “>” on the other.

Composition

Each tablet contains either 125, 250 or 500 mg of famciclovir as the active ingredient, and the following inactive ingredients: hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium starch glycolate, talc, and titanium dioxide.

Packaging

ACT FAMCICLOVIR tablets are available in HDPE bottles of 30, 100 and 500 tablets, and as aluminum foil-PVC/PVDC unit dose blister packages in the following pack sizes:

125 mg – 10 tablets per carton (1 strip of 10 tablets)

250 mg – 30 tablets per carton (3 strips of 10 tablets)

500 mg – 21 tablets per carton (3 strips of 7 tablets)


PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Common name:	Famciclovir
Chemical name:	2-[2-(2-amino-9H-purine-9-yl)-ethyl]-1,3-propanediol diacetate
Molecular formula:	C ₁₄ H ₁₉ N ₅ O ₄
Molecular mass:	321.34 g/mol

Structural Formula



Physicochemical properties:

Description:	White to off-white crystalline powder with a melting range of 101-105°C.
Solubility:	Soluble in water up to initial concentrations of > 25% w/v, but subsequently precipitates as the crystalline monohydrate. Solubility of the monohydrate in water is approximately 2.2% w/v at 25°C and approximately 9.1% w/v at 37°C.
pKa value:	3.84
pH value:	4.8 (1 mM solution)

CLINICAL TRIALS

Safety and Efficacy Trials

Herpes Zoster and Post-Herpetic Neuralgia

In patients with uncomplicated herpes zoster, famciclovir has been shown to significantly reduce the duration of virus shedding and to relieve the signs and symptoms of the disease.

A 7-day double-blind placebo-controlled trial was conducted in 419 patients with uncomplicated herpes zoster treated within 72 hours of initial lesion appearance. According to the randomization scheme, 138 patients were given famciclovir 500 mg t.i.d., 135 patients famciclovir 750 mg t.i.d. and 146 patients given placebo. No additional efficacy was demonstrated with the higher dose of famciclovir (750 mg t.i.d.), when compared to famciclovir 500 mg t.i.d. In the total population, 65.2% of patients had a positive viral culture at some time during their acute infection. Patients treated with famciclovir 500 mg had a shorter median duration of viral shedding (time to last positive viral culture) than did placebo-treated patients (1 day and 2 days, respectively; $p=0.0001$).

The times to loss of vesicles ($p=0.01$), loss of ulcers ($p=0.01$), and loss of crusts ($p=0.05$), were shorter for famciclovir 500 mg-treated patients than for placebo-treated patients in the overall study population.

The follow-up phase of this trial was designed to monitor the progression of post-herpetic neuralgia (PHN) after treatment with either famciclovir or placebo for seven days during acute infection. There was no difference in the incidence of postherpetic neuralgia between the treatment groups at the time of rash resolution. In the 186 patients (44.4% of total study population) who did develop postherpetic neuralgia, the median duration of postherpetic neuralgia was shorter in patients treated with famciclovir 500 mg than in those treated with placebo (63 days and 119 days, respectively; $p=0.02$).

A second 7-day double-blind controlled trial involved 545 patients with uncomplicated herpes zoster treated within 72 hours of initial lesion appearance. According to the randomization scheme, 134 patients received 250 mg t.i.d. of famciclovir, 134 patients received 500 mg t.i.d. of famciclovir, 138 patients received 750 mg t.i.d. of famciclovir and 139 patients received 800 mg of acyclovir given 5 times a day. Famciclovir was found to be as effective as acyclovir at all dose levels for cutaneous lesion healing parameters, time to loss of pain and viral shedding.

A double-blind controlled trial in 497 adult patients with ophthalmic zoster treated within 72 hours of initial lesion appearance compared famciclovir 500 mg three times daily for 7 days ($n=251$) to acyclovir 5 times per day for 7 days ($n=246$). Famciclovir was comparable to acyclovir in preventing ocular complications due to herpes zoster infection.

Herpes Simplex Infections

Treatment of Recurrent Genital Herpes Episodes:

Famciclovir was studied in two placebo-controlled trials of 626 otherwise healthy patients with a recurrence of genital herpes who were treated with famciclovir 125 mg b.i.d. (n=160), famciclovir 250 mg b.i.d. (n=169), famciclovir 500 mg b.i.d. (n=154) or placebo (n=143) for 5 days. In the two studies combined, the median time to healing in famciclovir 125 mg-treated patients was 4 days compared to 5 days in placebo treated patients (p= 0.0001) and the median time to cessation of viral shedding was 1.8 vs. 3.4 days in famciclovir 125 mg and placebo recipients, respectively (p= 0.0001). The median time to loss of all symptoms was 3.2 days in famciclovir 125 mg-treated patients vs. 3.8 days in placebo treated patients (p= 0.0001). Pre-treatment, self-obtained viral cultures were positive in 31%, 25%, 30% and 24% for the famciclovir 125 mg, 250 mg, 500 mg and placebo recipients respectively in the patient-initiated study. Of those patients whose pre-treatment culture was negative, significantly fewer patients self-initiating famciclovir treatment went on to become viral culture positive compared to placebo. Patients initiating treatment early (during the prodrome) were half as likely to commence viral shedding compared to placebo patients. Additionally, in the clinic-initiated study, famciclovir reduced the number of patients who developed new lesions.

Famciclovir was also studied in three acyclovir-controlled, double-blind trials in 951 otherwise healthy patients with first episode genital herpes.

Famciclovir for 5 or 10 days provided comparable efficacy to acyclovir although the studies were not powered to show statistical equivalence.

Suppression of Recurrent Genital Herpes Episodes:

A total of 934 otherwise healthy adults with frequently recurring genital herpes, were enrolled in two 12-month, placebo-controlled studies. Patients either had at least six recurrences 12 months prior to study entry or a history of at least six recurrences per year while not receiving other suppressive therapies. Sixty-two percent of patients had experienced at least 12 genital herpes recurrences in the previous 24 months. Treatment arms consisted of famciclovir 125 mg t.i.d. (n=233), 250 mg b.i.d. (n=236), 250 mg t.i.d. (n=232) and placebo (n=233). Compared to placebo, famciclovir 250 mg b.i.d. significantly delayed the time to developing the first clinically confirmed recurrence by 10 months in one study (medians: >365 days for famciclovir vs. 67 days for placebo; p=0.0001) and 9.5 months in another study (medians: 336 days for famciclovir vs. 47 days for placebo; p=0.0001). Approximately 80% of famciclovir-treated patients in both studies remained free from HSV recurrences documented by viral culture for up to 6 months compared with approximately 25% of patients treated with placebo (p<0.001). Treatment effects were sustained for 12 months.

Treatment of Recurrent Mucocutaneous Herpes Simplex Infection in HIV-Infected Patients:

A randomized, double-blind, multicenter study compared famciclovir 500 mg twice daily for 7 days (n=150) with oral acyclovir 400 mg 5 times daily for 7 days (n=143) in HIV-infected patients with mucocutaneous HSV infection. Approximately 40% of patients had a CD4 count below 200 cells/mm³; 54% of patients had anogenital lesions and 35% had orolabial lesions. Twice-daily oral famciclovir was comparable to five-times daily oral acyclovir in preventing new lesion formation, in time to complete healing (median of 7 days in both groups), time to loss of all lesion-associated symptoms (median of 4 days in both groups) and time to cessation of viral shedding (median of 2 days in both groups). Efficacy was maintained regardless of the degree of immunosuppression or location of lesions.

Comparative Bioavailability Studies

A blinded, single-dose, randomized, two-period, two-sequence, two-treatment crossover comparative bioavailability study of ACT FAMCICLOVIR (famciclovir) 500 mg tablets against the Canadian Reference Product, Famvir® (famciclovir) 500 mg tablets, has been performed on 28 healthy adult volunteers (male and female) under fasting conditions. A summary of the bioavailability data is presented in the table below.

Comparative Bioavailability Data for ACT FAMCICLOVIR (famciclovir) 500 mg vs. Famvir® (famciclovir) 500 mg Tablets

Analyte: Penciclovir (1 x 500 mg famciclovir) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	ACT FAMCICLOVIR * 500 mg tablets	Famvir® † 500 mg tablets	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (ng•h/mL)	11334.9 11606.1 (23)	10925.6 11120.9 (19)	103.75	100.74 - 106.84
AUC _I (ng•h/mL)	11653.1 11925.7 (22)	11230.3 11427.7 (19)	103.76	100.84 - 106.77
C _{max} (ng/mL)	4154.1 4259.6 (22)	3599.0 3681.8 (21)	115.43	107.36 - 124.09
T _{max} § (h)	0.79 (37)	0.93 (33)		
T _{1/2} § (h)	2.25 (13)	2.23 (16)		

* Famciclovir 500 mg Tablets (Actavis Pharma Company, Canada)

† Famvir® 500 mg Tablets (Novartis Pharmaceuticals Canada Inc., Canada)

§ Expressed as the arithmetic mean (CV%) only

VIROLOGY

Penciclovir is a highly potent and selective antiviral agent. Inhibitory activity in animals has been shown against herpes simplex virus type 1 and 2 (HSV-1 and HSV-2) infections. It also has inhibitory activity in cell cultures against HSV-1, HSV-2 and varicella zoster virus (VZV).

Penciclovir enters cells rapidly. In HSV-1, HSV-2 and VZV infected cells, viral thymidine kinase rapidly converts penciclovir to a monophosphate, which host cell enzymes convert to penciclovir triphosphate. The triphosphate inhibits viral DNA polymerase competitively with deoxyguanosine triphosphate and is incorporated into the extending DNA chain, preventing significant chain elongation. Consequently, viral DNA synthesis and, therefore, viral replication are inhibited.

Following the removal of the acyclonucleoside from MRC-5 cell culture medium, the stability of intracellular penciclovir triphosphate was much greater than that of acyclovir triphosphate. The intracellular half-life of penciclovir triphosphate was: 10 hours for HSV-1; 20 hours for HSV-2; 7 hours for VZV-infected cells grown in culture. The corresponding values for acyclovir triphosphate were 0.7 hours for HSV-1 and 1.0 hours for HSV-2. No value was derived for VZV-infected cells since acyclovir triphosphate concentrations were undetectable. In other human cells, penciclovir triphosphate was invariably a more stable entity than acyclovir triphosphate. The long intracellular half-life of penciclovir triphosphate ensures prolonged antiviral activity as demonstrated in cell cultures with HSV-1, HSV-2 and VZV, and in animal studies with HSV-1 and HSV-2.

In cell culture, penciclovir has the highest antiviral activity against the following herpes viruses (listed in decreasing order of potency and detailed below): HSV-1, HSV-2, VZV, Epstein-Barr virus (EBV) and cytomegalovirus (CMV). The degree of inhibition is dependent upon a number of variables, including the assay method, the host cell, virus type and multiplicity of infection. Comparative *in vitro* data for the antiviral potency of penciclovir and acyclovir are shown in the following table.

Assay Method	Virus*	Cell Type	Penciclovir	Acyclovir
IC50 (mcg/mL)				
Plaque reduction	VZV	MRC-5	2.0 – 10.9	2.8 – 8.5
	VZV	Hs68	0.4 – 1.8	0.4 – 1.6
	HSV-1	WISH	0.04 – 0.5	0.2 – 1.2
	HSV-1	MRC-5	0.2 – 0.6	0.1 – 0.6
	HSV-2	WISH	0.1 – 0.8	0.4 – 1.8
	HSV-2	MRC-5	0.9 – 2.1	0.2 – 1.1
DNA synthesis inhibition	VZV (Ellen)	MRC-5	0.1	0.2
	HSV-1 (SC-16)	MRC-5	0.04	0.15
	HSV-2 (MS)	MRC-5	0.05	0.04
IC99 (mcg/mL)				
Virus yield reduction	HSV-1	MRC-5	0.4 – 0.6	0.8 – 2.5
	HSV-2	MRC-5	0.6 – 0.7	0.2 – 0.9

* Data were obtained using clinical isolates, except for DNA synthesis assays in which the specified laboratory strains were used.

In an assay measuring inhibition of EBV DNA synthesis in P₃HR1 cells, the IC₅₀ for penciclovir was 1.5 mcg/mL. In plaque-reduction assays in MRC-5 cells against CMV, the IC₅₀ for penciclovir was 52 mcg/mL.

Like with aciclovir, penciclovir resistance is associated with mutations principally in the TK gene resulting in deficiency or altered substrate specificity of this enzyme, and to a much lesser extent in the DNA polymerase gene. Most aciclovir-resistant HSV and VZV clinical isolates are also resistant to penciclovir, but cross-resistance is not universal. In plaque-reduction assays against laboratory-produced mutants with altered DNA polymerase, Vero (human lung fibroblast) cells infected with HSV-1 were inhibited by penciclovir to the same extent as the wild type (IC₅₀ was 1.5 mcg/mL for the mutant and 1.0 mcg/mL for the wild type). In contrast, resistance to acyclovir was demonstrated as the IC₅₀ of acyclovir was 47 fold higher (20 mcg/mL) for the mutant compared to (0.43 mcg/mL) the wild type. In a clinically isolated strain of acyclovir-resistant HSV-1 tested in Hs68 (human foreskin fibroblast) cells, resistance to acyclovir was demonstrated, whereas the IC₅₀ values for penciclovir in the wild type and mutant strains were almost identical (1.3 mcg/mL vs. 1.0 mcg/mL, respectively).

Penciclovir sensitivity testing was performed on 84 HSV-2 isolates from 50 famciclovir-treated patients (59 isolates) and 21 placebo-treated patients (25 isolates) who participated in a 4-month genital herpes suppression trial. All HSV-2 isolates on-treatment and after-treatment were sensitive (mean IC₅₀ 1.7 mcg/mL for famciclovir recipients and 1.5 mcg/mL for placebo recipients).

Short-term treatment of HSV-1-infected MRC-5 cells with penciclovir (3 mM) for 2 hours reduced viral DNA by 76% compared with only 17% following similar treatment with acyclovir (3 mM). Treatment of HSV-2-infected cells with penciclovir (3 mM) for 1.5 hours reduced viral DNA by 52% vs. 20% with acyclovir (3 mM).

Studies of VZV-infected MRC-5 cells exposed to pulse treatment for 8 hours on days 0, 1, 2 and 3 produced an IC₅₀ of 5.0 mcg/mL for penciclovir and 24 mcg/mL for acyclovir. The activity of penciclovir seen after short or pulse treatment is consistent with the stability of intracellular penciclovir triphosphate.

The presence of penciclovir at 10 or 100 mcg/mL did not decrease the activity of zidovudine against human immunodeficiency virus in M8166 cells.

As there is no appropriate animal model that mimics VZV infection in humans, the antiviral activity of penciclovir has not been evaluated in animals infected with VZV. However, penciclovir has been shown to have inhibitory activity against HSV-1 and HSV-2 infections in mice and guinea pigs. The level of antiviral activity depends on a number of factors including the route of infection, route of administration of penciclovir and time between virus infection and treatment with penciclovir.

The clinical significance of inhibitory activity of penciclovir against HSV-1 and HSV-2 in *in vitro* animal models is unknown at this time.

TOXICOLOGY

Both famciclovir and its active metabolite penciclovir have undergone a comprehensive toxicological evaluation; principal findings from the key studies are summarised below. Exposure to drug-related material in animals compared with that in humans given 500 mg famciclovir t.i.d. is expressed, where appropriate, as multiples of the amount of penciclovir and 6-deoxy-penciclovir systemically available.

Acute Toxicity:

Famciclovir has a low order of acute toxicity, with median lethal doses in excess of 5000 mg/kg orally and around 500 mg/kg or higher intravenously; penciclovir intravenously was slightly better tolerated. With both compounds the majority of adverse effects were indicative of central nervous system disturbances, undoubtedly reflecting the high doses administered since no such effects were detected at the lower doses used in the studies to assess general pharmacological actions.

Subacute Toxicity Studies:

In studies of 1 month's duration, famciclovir was administered at oral doses of up to 4000 mg/kg/day in the rat and up to 500 mg/kg/day in the dog.

In rats, there were early degenerative changes in the seminiferous epithelium at 4000 and 400 mg/kg/day (35- and 9- fold human exposure to penciclovir respectively), but the effects at the lower dose were minimal. Microscopically there was also a slight reduction in the number of lymphocytes in the thymic cortex confined to the high dose of 4000 mg/kg/day; there were no effects at 400 mg/kg/day.

In dogs, doses of up to 500 mg/kg/day (3-fold human exposure to penciclovir), which was the maximum practicable dose, were tolerated with no indications of definitive target organ toxicity.

Juvenile toxicity study in rats

In juvenile rats, famciclovir was administered daily at doses of 0, 40, 125, or 400 mg/kg/day for 10 weeks beginning on post-partum Day 4. There were no treatment related deaths clinical observations or adverse developmental effects. The toxicity of famciclovir was not enhanced in juvenile rats compared to that in the adult animals.

Chronic Toxicity Studies:

In 6 and 12 month studies with famciclovir, oral doses of up to 600 mg/kg/day in rats and up to 500 mg/kg/day in dogs were without significant effect on any tissue except the testes (See

Reproduction). At these doses, a high systemic exposure to both penciclovir (4- and 3-fold human values in rat and dog respectively) and 6-deoxy penciclovir (at least 40- and 150-fold human values in rat and dog respectively) were achieved.

Carcinogenicity Studies:

Two-year dietary carcinogenicity studies with famciclovir were conducted in rats and mice. There was an increase in the incidence of mammary adenocarcinoma in female rats receiving the maximum tolerated dose of 600 mg/kg/day (4- and 40-fold human exposure to penciclovir and 6-deoxy penciclovir respectively), but not in the number of high dose females with mammary tumours of any type (benign or malignant). No effects were seen in females receiving 200 mg/kg/day (2- and 8-fold human exposure to penciclovir and 6-deoxy penciclovir respectively) or in males treated at doses up to 240 mg/kg/day, the maximum tolerated dose. There was also no effect on the incidence or type of tumours in mice given the maximum tolerated dose of 600 mg/kg/day (6- and 65-fold human exposure to penciclovir and 6-deoxy penciclovir respectively).

Mutagenicity Studies:

Famciclovir was not found to be genotoxic in a comprehensive battery of *in vivo* and *in vitro* tests designed to detect gene mutation, chromosomal damage and repairable damage to DNA. Penciclovir, in common with other drugs of this class, has been shown to cause chromosomal damage, but did not induce gene mutation in bacterial or mammalian cell system. There was also no evidence of increased DNA repair *in vitro*. Penciclovir caused an increased incidence of micronuclei in mouse bone marrow *in vivo* when administered intravenously at doses highly toxic to bone marrow (500 mg/kg and above) but not when administered orally at 2,400 and 4,800 mg/kg.

Reproduction:

As with other drugs of this class, testicular toxicity has been observed in animals receiving both famciclovir and penciclovir, but there was evidence of a return towards normal on withdrawal of treatment. The testicular effects seen following famciclovir are very likely attributable to penciclovir; therefore, systemic exposure comparisons in animals and man are shown for penciclovir.

Mating performance was unaffected in male rats treated with famciclovir for up to 6 months, but fertility decreased after 10 weeks of dosing at 500 mg/kg/day (5-fold human exposure) and there were abnormal sperm profiles, degenerative changes in the seminiferous epithelium and raised FSH levels. Fertility was unaffected at 150 mg/kg/day (2-fold human exposure).

Only minimal testicular degeneration was apparent in rats after 1 month's dosing at 400 mg/kg/day (9- fold human exposure), and no effects were seen following 240 mg/kg/day for 2 years (3-fold human exposure).

In dogs there were no effects after 1 month's dosing at 500 mg/kg/day (3-fold human exposure) and, except for minor histological changes in a single dog after 12 months' treatment, testicular toxicity was confined to chronic dosing with ≥ 150 mg/kg/day.

There were no significant effects on fertility or on peri- and post-natal development in female rats given famciclovir at doses up to 1000 mg/kg/day.

Famciclovir was tested for effects on embryo-fetal development in rats and rabbits at oral doses up to 1000 mg/kg/day and intravenous doses of 360 mg/kg/day in rats or 120 mg/kg/day in rabbits. No adverse effects were observed on embryo-fetal development. Similarly, no embryotoxic or teratogenic effects were observed following intravenous administration of penciclovir to rats (up to 80 mg/kg/day) or rabbits (up to 60 mg/kg/day).

There were no clinically significant effects on sperm count, morphology or motility in male patients receiving famciclovir 250 mg b.i.d. for 18 weeks. Although the daily dosage in this study was lower than the recommended dose for zoster, the duration of treatment was longer at 18 weeks compared with 7 days, and patients therefore received a cumulative (total) dose which was 6-fold higher.

REFERENCES

1. Boyd M.R., Safrin S., Kern E.R. Penciclovir: A Review of its Spectrum of Activity, Selectivity, and Cross-resistance pattern. *Antiviral Chemistry and Chemotherapy* 1993; 4 (Suppl. 1): 3-11
2. Daniels S., Schentag J.J. Drug Interaction Studies and Safety of Famciclovir in Healthy Volunteers: A Review. *Antiviral Chemistry and Chemotherapy* 1993; 4 (Suppl. 1): 57-64
3. Degreef H. et al. (Famciclovir Herpes Zoster Clinical Study Group). Famciclovir, a New Oral Antiherpes Drug: Results of the First Controlled Clinical Study Demonstrating its Efficacy and Safety in the Treatment of Uncomplicated Herpes Zoster in Immunocompetent Patients. *International Journal of Antimicrobial Agents* 1994; 4 (4): 241-246
4. Portnoy J. Famciclovir in the Treatment of Herpes Zoster (HZ) Infection. *Antiviral Research* 1994; 23(Suppl. 1): 98
5. Pue M.A., Benet L.Z. Pharmacokinetics of Famciclovir in Man. *Antiviral Chemistry and Chemotherapy* 1993; 4 (Suppl. 1): 47-55
6. Saltzman R., Jurewicz R., Boon R. Safety of Famciclovir in Patients with Herpes Zoster and Genital Herpes. *Antimicrobial Agents and Chemotherapy* 1994; 38(10): 2454-2457
7. Tyring S., Barbarash R., Nahlik J., Cunningham A., Marley J., Heng M., Jones T. Efficacy of Famciclovir on Herpes Zoster Rash Resolution and Post-herpetic Neuralgia. *Antiviral Research* 1994; 23(Suppl. 1): 73
8. Tyring S., Barbarash R., Nahlik J. et al. Famciclovir for the Treatment of Acute Herpes Zoster: Effects on Acute Disease and Postherpetic Neuralgia. *Ann Int Med* 1995; 123:89-96.
9. Vere Hodge R.A., Cheng Y-C. The Mode of Action of Penciclovir. *Antiviral Chemistry and Chemotherapy* 1993; 4 (Suppl. 1): 13-24
10. Whitley R.J. Penciclovir/Famciclovir: Parent and Child in the Treatment of Herpes Virus Infections. *Expert Opinion on Investigational Drugs* 1994; 3 (7): 759-761
11. Sacks S, Aoki F, Diaz-Mitoma F, Sellors J, Shafran S et al. Patient-Initiated, Twice-Daily Oral Famciclovir for Early Recurrent Genital Herpes A Randomized, Double-blind Multicentre Trial. *JAMA* 1996; 276: 44-49.
12. Perry CM, Wagstaff AJ. Famciclovir - A Review of its Pharmacological Properties and Therapeutic Efficacy in HerpesVirus Infections. *Drugs* 1995; 50: 396-415.
13. Thackray AM, Field H. Differential Effects of Famciclovir and Valacyclovir on the Pathogenesis of Herpes Simplex Virus in a Murine Infection Model Including Reactivation from Latency. *Journal of Infectious Diseases* 1996; 173: 291-299
14. Thackray AM, Field H. Comparison of Effects of Famciclovir and Valaciclovir on Pathogenesis of Herpes Simplex Virus Type 2 in a Murine Infection Model. *Antimicrobial Agents and Chemotherapy* 1996; 40: 846-851.
15. Sacks S, Martel A, Aoki F, Shafran S, St-Pierre C, Lassonde M et al. Early Clinic-initiated Treatment of Recurrent Genital Herpes using Famciclovir, Results of a Canadian Multicentre Study. *Clin Res* 1994; 42: 300A.
16. Sacks SL, Aoki FY, Diaz-Mitoma F, Sellors J, Shafran S, Romanowski B, Martel A,

- Williams K, Haase D et al. Patient and Clinic-initiated Treatment of Recurrent Genital Herpes with Twice-Daily Oral Famciclovir. 7th European Congress of Clinical Microbiology and Infectious Diseases 1995; 866A.
17. Loveless M, Harris W, Sacks S. Treatment of First Episode Genital Herpes with Famciclovir. 35th ICAAC 1995; Abstract H12.
 18. Frechette G, Romanowski B. Efficacy and safety of famciclovir for the treatment of HSV infection in HIV+ patients. Can J Infect Dis 1997; 8(suppl A): 44A: 301.
 19. Product Monograph for Famvir[®] (famciclovir). Novartis Pharmaceuticals Canada Inc., Quebec. Control Number: 196268; Date of Revision: September 14, 2016.

PART III: CONSUMER INFORMATION

PrACT FAMCICLOVIR (famciclovir)

This leaflet is part III of a three-part "Product Monograph" published when ACT FAMCICLOVIR (famciclovir) was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about ACT FAMCICLOVIR. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

ACT FAMCICLOVIR is an antiviral medicine that is used to treat a number of viral infections described below. It stops the infecting virus from reproducing. Since the virus reproduces very early in the infection, the best results in the treatment are obtained if ACT FAMCICLOVIR is started as soon as possible after the first symptoms appear.

Herpes zoster:

ACT FAMCICLOVIR is used to treat herpes zoster (Shingles). Herpes zoster is an infection caused by a virus called varicella zoster (the same virus that causes chickenpox). ACT FAMCICLOVIR stops the virus from spreading in the body so that healing can occur faster if taken within 72 hours of the first lesion, and reduces the blistering, pain, and rash.

Genital herpes:

ACT FAMCICLOVIR is used to treat or prevent recurrence of the viral infections which cause genital herpes. Genital herpes is a viral infection caused by herpes simplex virus type 1 or 2, and is normally spread by sexual contact. It causes blisters and burning or itching around the genitals, which may be painful.

ACT FAMCICLOVIR does not stop you from spreading herpes to another person. It is important to understand that the drug must be taken as early as possible as soon as you know that an attack is starting. Although ACT FAMCICLOVIR does not cure the viral infection, it helps to shorten the duration of symptoms and allows the infections to heal faster.

What it does:

ACT FAMCICLOVIR helps stop the virus from multiplying and shortens the time the virus is released and spread, which helps healing.

When it should not be used:

Do not use ACT FAMCICLOVIR if:

- you have a known hypersensitivity to famciclovir, to

penciclovir (a compound formed from famciclovir by your body and an ingredient of some other medicines) (See "What the important nonmedicinal ingredients are) or to any ingredient of the formulation or packaging

What the medicinal ingredient is:

Each ACT FAMCICLOVIR tablet contains 125mg, 250 mg or 500 mg of famciclovir.

What the important nonmedicinal ingredients are:

The tablets also contain some inactive ingredients to make up the bulk of each tablet: Hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium starch glycolate, talc, and titanium dioxide.

The tablets are gluten-free.

The tablets do not contain sucrose, tartrazine or any other azo dyes.

What dosage form it comes in:

ACT FAMCICLOVIR is available as 125 mg, 250 mg or 500 mg tablets with the following descriptions:

ACT FAMCICLOVIR 125 mg tablets are white to off-white, film-coated, round biconvex tablet with "FC" on one side and "S" on the other. 125

ACT FAMCICLOVIR 250 mg tablets are white to off-white, film-coated, round biconvex tablet with "FC" on one side and "S" on the other. 250

ACT FAMCICLOVIR 500 mg tablets are white to off-white, film-coated, capsule-shaped, biconvex tablet with "FC500" on one side and "S" on the other.

Check with your doctor or pharmacist if the identifying markings or colour of your tablets are not the same as these.

WARNINGS AND PRECAUTIONS

Before you take ACT FAMCICLOVIR, **talk to your doctor if:**

- You have an allergy (including itching) or other unwanted (side) effects when previously taking famciclovir
- You are pregnant, planning to become pregnant, breastfeeding or planning to breastfeed
- You have a past history or current kidney disease
- You have severe liver problems
- You have other medical conditions

It is important to tell your doctor, dentist or pharmacist if you are taking other medication, as combining drugs can sometimes result in a change from the expected drug effects, or cause harmful effects.

INTERACTIONS WITH THIS MEDICATION

Always tell your doctor about other medical conditions you have or any medicines you are taking. This means medicines you bought for yourself as well as medicines on prescription. Your doctor or pharmacist will know if it is safe to take ACT FAMCICLOVIR as well.

It is particularly important that you tell your doctor or pharmacist if you are taking any of the following medicines:

- Probenecid (used to treat high blood levels of uric acid associated with gout and to increase blood levels of penicillin-type antibiotics), or any other medicine that can affect your kidneys.
- Raloxifene (used to prevent and treat osteoporosis).

You can take ACT FAMCICLOVIR with or without food.

PROPER USE OF THIS MEDICATION

How do I use ACT FAMCICLOVIR?

You should take ACT FAMCICLOVIR only as directed by your doctor. Do not take more of it, do not take it more often and do not take it for a longer period of time than your doctor ordered.

Be sure to take ACT FAMCICLOVIR regularly as prescribed. Try to take your tablets at the same time each day. You should continue to take your medicine even if you do not feel better, as it may take a few days for your medicine to work.

If your kidneys are not working very well, your doctor will probably have you take this medicine less often, check with your doctor if you have kidney problems.

ACT FAMCICLOVIR will work whether or not you take it with food. You should swallow the tablets whole, with water. Do not chew them.

How long do I use ACT FAMCICLOVIR?

The usual adult dose is one ACT FAMCICLOVIR tablet three times a day for shingles. Follow your doctor's instructions about how and when to take your tablets. Start taking them as soon as you can for the best effect.

Most people take one tablet when they get up in the morning, one in the middle of the afternoon and one before going to bed at night.

Finish all the tablets you have been given even if you start to feel better. The treatment lasts for seven days.

Treating an outbreak of recurrent genital herpes:

If you have had herpes infections of the genitals before (called recurrent genital herpes), your doctor may decide to treat the outbreak. You will likely be told to take one 125 mg tablet twice a day for 5 days or if you are HIV-positive, one 500 mg tablet twice a day for 7 days. Most people with recurrent genital herpes take one tablet when waking, and one tablet just before going to bed. ACT FAMCICLOVIR should be taken as soon as possible after the early symptoms (pain, burning, blisters, itching, tingling) begin to appear.

Preventing outbreaks of recurrent genital herpes:

If you have had herpes infections of the genitals before (called recurrent genital herpes), then your doctor may decide to prevent future outbreaks. Your doctor will likely tell you that you should continually take one 250 mg tablet twice a day. Most people with recurrent genital herpes take one tablet when waking, and one tablet just before going to bed.

Missed Dose:

If you miss taking a ACT FAMCICLOVIR tablet, don't worry. Take it as soon as you remember. Take your next tablet at the normal time. However, do not take two doses within a time interval of less than 1 hour, in that case you should skip the missed dose. Carry on as normal until you have finished all the tablets. It is important that you finish all the tablets you have been given, unless your doctor tells you to stop taking them. Do not double-dose.

Overdose

In case of drug overdose, contact a healthcare practitioner (doctor), hospital emergency department or regional poison control center, even if there are no symptoms. Bring your pack of tablets with you.

Taking too much of any type of medicine is risky.

Taking too much ACT FAMCICLOVIR may affect the kidneys. In people who already have kidney problems it may, rarely lead to kidney failure if their dose is not correctly lowered.

Important: Your doctor may give you different instructions better suited to your specific needs. If you need more information on how to take ACT FAMCICLOVIR properly, double-check with your doctor or pharmacist.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Any medicine can cause side effects. With ACT FAMCICLOVIR, some people may feel sick or get a headache, or experience confusion or dizziness. Other side effects include drowsiness (usually in older people), hallucinations (seeing or hearing things that are not really there), vomiting, yellowing of the skin and/or eyes, abnormal liver function test results, palpitations (signs of abnormal heart beat), skin rash, itching or blistering, diarrhea, tiredness and abdominal pain. If you get

these or any other problems while you are taking ACT FAMCICLOVIR, tell your doctor or pharmacist. Some effects could be serious:

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM			
Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Signs of serious skin reaction: pruritus, (e.g. itching, erythema multiforme, Steven Johnson syndrome, toxic epidermal necrolysis), severe blistering of the skin and/or mucous membranes of the lips, eyes, mouth, nasal passages or genitals, angioedema (e.g. swelling of tissues such as face, mouth, eyes, throat, skin, hands and feet)			√
Serious allergic reaction with symptoms such as difficulty of breathing or swallowing, rash, itching, hives, wheezing or coughing, light-headedness, dizziness, changes in levels of consciousness, hypotension, with or without generalized itching, skin reddening, facial/throat swelling, blue discoloration of the lips, tongue or skin			√
Seizures or fits			√
Signs of decrease in the number of blood platelets: Unexplained bruising, reddish or purplish patches on the skin or nosebleeds			√
Purple skin patches, itching, burning (signs of inflamed blood vessels)			√

Tell your doctor immediately or go to the emergency department at your nearest hospital if any of the following side effects occur.

If you experience anything unusual, while you are receiving treatment with ACT FAMCICLOVIR, you should report it to your physician.

A more complete listing of side effects reported to date, is contained in the Product Monograph supplied to your physician.

Driving and using machines

ACT FAMCICLOVIR can cause dizziness, drowsiness or confusion in very rare cases. If you have any of these symptoms while taking ACT FAMCICLOVIR, you should take special care when driving or using machines.

This is not a complete list of side effects. For any unexpected effects while taking ACT FAMCICLOVIR, contact your doctor or pharmacist.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to: 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Heath Canada
Postal Locator 1908C
Ottawa ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Website at www.healthcanada.gc.ca/medeffect.

NOTE: *Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice."*

HOW TO STORE IT

- Keep ACT FAMCICLOVIR in the bottle or package provided by the pharmacist.
- Store at room temperature (15-30°C).
- Protect from light.
- ACT FAMCICLOVIR must be kept out of the reach and sight of children.
- Don't take your tablets if they are past their expiry date.
- Never let anyone else take your tablets, even if they have shingles as well.
- Finish all the tablets in the way you have been told.

More information about shingles.

What is shingles like?

- At first you might feel burning and tingly where the rash is coming. You may get pain for a few days before you see the rash.
- Most people with shingles get a blistery rash down one side of their body or face. This can be painful.
- New blisters will keep coming for about five days. After this, the blisters will dry to form scabs.
- You may feel weak and tired.
- The rash usually lasts for two to three weeks. Afterwards, people can have pain where the rash was, sometimes for several months.

Who gets shingles?

People of any age can get shingles. Most people who get it are middle-aged or older.

You can only get shingles if you have had chickenpox.

50% of people have had shingles by the age of 85.

What causes shingles?

- Shingles is caused by the same virus that causes chickenpox.
- After you have had chickenpox, the inactive virus stays in your body.
- Many years later the virus can start to work again. This may happen when you are run down or tired.

How is shingles treated?

Tablets for shingles, such as ACT FAMCICLOVIR, stop the virus spreading. They cut down blistering and pain of the rash. They help you get better quicker if you take them early in your illness.

What else can you do?

- To stop the rash itching, have cool baths - do not use perfumed soap or bath oils. You can also put ice cubes wrapped in a washcloth on the rash, or cover it with a soothing lotion like calamine for the first three days.
- Keep the rash clean and dry.
- Wear loose clothes.
- Do not scratch the rash. It could get infected and take longer to get better.
- Rest when you feel tired.
- Try to eat well and drink plenty of fluids.
- Tell your doctor right away if you start to have trouble with your eyes. Shingles can sometimes cause eye problems which can be treated.

Can other people catch your shingles?

Shingles is caused by the same virus that causes chickenpox. Most doctors agree that people do not catch shingles from each other. When you have shingles, you can give chickenpox to someone who has never had chickenpox before - but this is unlikely.

What is Genital Herpes?

Genital Herpes is a viral infection of the genital (sex organ) area which is caused by the Herpes Simplex Virus. You might suspect the onset of this infection if you see the appearance of sores or blisters or feel a burning sensation in your genital region.

Type II herpes simplex virus is the main cause of the sores and blisters that develop in your genital area, but you can also catch genital herpes from herpes simplex Type I which can be the cause of cold sores which occur around your mouth.

Genital Herpes can occur as a first-episode attack or recurrent infection. Unlike many other viruses, recovery from the first attack does not leave you with lifelong protection from re-infection. The virus is able to remain hidden in the nerves after the initial infection and stays there until reactivated.

Since your body still carries the highly infectious virus, you can easily infect someone else, even if you feel fine and you have no symptoms of genital herpes. This explains why genital herpes is one of the most common sexually transmitted diseases (STD).

The risk is higher for people who are more sexually active and have more sexual partners. It is therefore recommended that you avoid sexual activity if you or your partner have any symptoms of herpes, even if you have started your treatment.

"If you are taking ACT FAMCICLOVIR to treat or to suppress genital herpes, or you have had genital herpes in the past, you should still practice "safer sex", including the use of condoms. This is important to prevent you passing the infection on to others.

While taking this medication:

- remember that your infection is contagious.
- tell any other doctor, pharmacist or dentist you see, that you are taking ACT FAMCICLOVIR.
- contact your doctor if you develop any unusual discomfort.
- you should not take ACT FAMCICLOVIR when pregnant or breast-feeding unless your doctor tells you to.
- do not give ACT FAMCICLOVIR to others because it may not be suitable for them.
- store your tablets in a dry place at room temperature in the original container provided by the pharmacy.
- keep this medication out of reach of children.
- read your prescription label carefully; consult your doctor or pharmacist if you have any questions or require further information.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting the sponsor, Actavis Pharma Company, at: 1-866-254-6111.

This leaflet was prepared by:
Actavis Pharma Company

6733 Mississauga Road, Suite 400
Mississauga, ON
Canada, L5H 6J5

Last revised: December 06, 2016