

PRODUCT MONOGRAPH

^{Pr}TEVA-MOMETASONE

Mometasone Furoate 0.1% Ointment, USP

Topical Corticosteroid

Teva Canada Limited
30 Novopharm Court
Toronto, Canada
M1B 2K9
www.tevacanada.com

Date of Revision:
May 7, 2018

Control # 215362

Table of Contents

| | |
|--|-----------|
| PART I: HEALTH PROFESSIONAL INFORMATION | 3 |
| SUMMARY PRODUCT INFORMATION | 3 |
| INDICATIONS AND CLINICAL USE | 3 |
| CONTRAINDICATIONS | 3 |
| WARNINGS AND PRECAUTIONS | 4 |
| ADVERSE REACTIONS | 7 |
| DRUG INTERACTIONS | 9 |
| DOSAGE AND ADMINISTRATION | 9 |
| OVERDOSAGE | 10 |
| ACTION AND CLINICAL PHARMACOLOGY | 11 |
| STORAGE AND STABILITY | 11 |
| DOSAGE FORMS, COMPOSITION AND PACKAGING | 11 |
| PART II: SCIENTIFIC INFORMATION..... | 12 |
| PHARMACEUTICAL INFORMATION | 12 |
| CLINICAL TRIALS | 13 |
| DETAILED PHARMACOLOGY | 17 |
| TOXICOLOGY | 18 |
| REFERENCES | 20 |
| PART III: CONSUMER INFORMATION | 25 |

TEVA-MOMETASONE

Mometasone furoate

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

| Route of Administration | Dosage Form / Strength | Clinically Relevant Nonmedicinal Ingredients |
|-------------------------|------------------------|--|
| Topical | Ointment / 0.1% | <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i> |

INDICATIONS AND CLINICAL USE

TEVA-MOMETASONE (mometasone furoate):

- is indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses such as psoriasis and atopic dermatitis.
- can be used for a maximum of 3 weeks duration on the body and of 5 days duration on the face, scalp, axillae and scrotum.

Geriatrics (≥ 65 years of age): No overall differences in safety and effectiveness were observed between subjects greater than 65 years of age and younger subjects receiving mometasone ointment. Geriatric patients may be more susceptible to the potential effects of systemic absorption. Decreased hepatic or renal function in the elderly may delay elimination.

Pediatrics (<18 years of age): Safety and efficacy of mometasone have not been established in pediatric patients.

CONTRAINDICATIONS

TEVA-MOMETASONE (mometasone furoate) is contraindicated:

- in patients who are sensitive to mometasone furoate, to other corticosteroids or to any component of these preparations. For a complete listing, see the Dosage Forms, Composition, and Packaging section of the product monograph.
- In patients with viral (e.g. herpes or varicella) lesions of the skin, bacterial or fungal skin infections, parasitic infections, skin manifestations relating to tuberculosis or syphilis, eruptions following vaccinations, acne vulgaris, rosacea, pruritus without inflammation.

- For use in the eyes.
- For use with occlusive dressings

WARNINGS AND PRECAUTIONS

General

Patients should be advised to inform subsequent physicians of the prior use of corticosteroids.

TEVA-MOMETASONE should not be used under occlusion, due to increased risk of systemic exposure and infection. When used under occlusive dressing, which is contraindicated, over extensive areas or on the face, scalp, axillae or scrotum, sufficient absorption may occur to result in adrenal suppression and other systemic effects (see WARNINGS AND PRECAUTIONS — Endocrine and Metabolism, Immune and Ophthalmologic).

Carcinogenesis and Mutagenesis

In genetic toxicity studies, mometasone furoate was not mutagenic in bacteria (Ames test) or mammalian (mouse lymphoma) cells and was not clastogenic in the mouse micronucleus test.

Cardiovascular

Suitable precautions should be taken when using topical corticosteroids in patients with stasis dermatitis and other skin diseases with impaired circulation.

Use of corticosteroids around chronic leg ulcers may be associated with a higher occurrence of local hypersensitivity reactions and an increased risk of local infection.

Endocrine and Metabolism

Manifestations of hypercortisolism (Cushing's syndrome) and reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, leading to glucocorticosteroid insufficiency, can occur in some individuals as a result of increased systemic absorption of topical corticosteroids. Hyperglycemia and glucosuria may occur in some patients due to systemic absorption of topical corticosteroids (see ADVERSE REACTIONS).

Conditions which augment systemic absorption include the formulation and potency of the topical corticosteroid, the application of topical corticosteroids over large body surface areas, application to intertriginous areas (such as the axillae), frequency of application, prolonged use or the addition of occlusive dressings. Other risk factors for increased systemic effects include increasing hydration of the stratum corneum,

use on thin skin areas (such as the face), use on broken skin or conditions where the skin barrier may be impaired.

If patients must be treated over large body surface areas, they should be evaluated periodically for evidence of HPA axis suppression (see WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests). If HPA axis suppression or Cushing’s syndrome is observed, an attempt should be made to withdraw the drug by reducing the frequency of the application. Abrupt withdrawal of treatment may result in glucocorticosteroid insufficiency (see ADVERSE REACTIONS).

Recovery of HPA axis function is generally prompt upon discontinuation of topical corticosteroids. Infrequently, signs and symptoms of glucocorticosteroid insufficiency may occur requiring supplemental systemic corticosteroids. For information on systemic corticosteroid supplementation, see prescribing information for those products.

Pediatric patients may absorb larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity from equivalent doses because of their larger skin surface to body mass ratios as compared with adults (see WARNINGS AND PRECAUTIONS – Special Populations, Pediatrics).

Immune

Topical corticosteroids may increase the risk of infections including aggravation of cutaneous infection, masked infection and secondary infections. In particular, bacterial infection is encouraged by warm, moist conditions within skin-fold areas or caused by occlusive dressings. If concomitant skin infections develop, TEVA-MOMETASONE should be discontinued and antimicrobial therapy administered.

Ophthalmologic

Topical corticosteroids should be used with caution on lesions close to the eye because systemic absorption may cause increased intraocular pressure, glaucoma or cataracts.

Visual disturbance may be reported with systemic and topical (including, intranasal, inhaled and intraocular) corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes of visual disturbances; this may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

Sensitivity

Local hypersensitivity reactions (see ADVERSE REACTIONS) may resemble symptoms of the condition under treatment. If hypersensitivity reactions occur, the drug should be discontinued and appropriate therapy initiated.

Allergic contact dermatitis with corticosteroids is usually diagnosed by observing a failure to heal rather than noticing a clinical exacerbation. Such an observation

should be corroborated with appropriate diagnostic patch testing.

Sexual Function/ Reproduction

There are no data in humans to evaluate the effect of topical corticosteroids on fertility.

Skin

If irritation or sensitization develops with the use of TEVA-MOMETASONE products, treatment should be discontinued and appropriate therapy instituted.

Prolonged use of topical corticosteroid preparations may produce striae or atrophy of the skin or subcutaneous tissue. Topical corticosteroids should be used with caution on lesions of the face, groin and axillae as these areas are more prone to atrophic changes than other areas of the body. Frequent observation is important if these areas are to be treated. If skin atrophy is observed, treatment should be discontinued.

Special Populations

Pregnant Women: The safe use of topical corticosteroids during pregnancy has not been established. Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at dosage levels that are similar to therapeutic doses. Some corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. The relevance of this finding to human beings has not been established.

In reproduction studies conducted in rats and rabbits, mometasone furoate produced effects such as reduced maternal body weight gain, suppression of fetal growth, delayed ossification, umbilical hernias, prolonged gestation, difficult and prolonged labor and inability to deliver (see TOXICOLOGY).

TEVA-MOMETASONE should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. The minimum quantity should be used for the minimum duration.

Nursing Women: Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. The safe use of topical corticosteroids during lactation has not been established. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when TEVA-MOMETASONE is administered to a nursing woman. Administration of TEVA-MOMETASONE during lactation should only be considered if the expected benefit to the mother outweighs the risk to the infant. If used during lactation, TEVA-MOMETASONE should not be applied to the breasts to avoid accidental ingestion by the infant.

Pediatrics (< 18 years of age): Safety and efficacy of mometasone have not been

established in pediatric patients.

Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA axis suppression and Cushing's syndrome when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency during and/or after withdrawal of treatment.

Adverse effects including striae have been reported with use of topical corticosteroids in infants and children. HPA axis suppression, Cushing's syndrome, linear growth retardation, delayed weight gain, and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include low plasma cortisol levels and an absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema. Chronic corticosteroid therapy may interfere with the growth and development of children.

Geriatrics (≥ 65 years of age): No overall differences in safety were observed between subjects greater than 65 years of age and younger subjects receiving TEVA-MOMETASONE. Topical corticosteroids should be used cautiously in elderly patients, reflecting their increased skin fragility and greater frequency of hepatic, renal, or cardiac dysfunction, and of concomitant disease or other drug therapy. Decreased hepatic or renal function in the elderly may delay elimination if systemic absorption occurs. The minimum quantity should be used for the minimum duration.

Patients with renal/hepatic impairment: Safety of mometasone has not been established in patients with renal or hepatic impairment. In case of systemic absorption, metabolism and elimination may be delayed leading to increased risk of systemic toxicity; therefore, minimum quantity should be used for the minimum duration.

Monitoring and Laboratory Tests

The cosyntropin (ACTH₁₋₂₄) stimulation test may be helpful in evaluating patients for HPA axis suppression.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Local adverse reactions rarely reported with mometasone ointment 0.1% include burning, pruritus, tingling/stinging and signs of skin atrophy. In <1% of patients, adverse reactions reported with mometasone ointment 0.1% include aggravated allergy, dermatitis, erythema, furunculosis, increased lesion size, nausea (one patient) and vaginal discharge (one patient).

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

The overall incidence of side effects was 4.9%, i.e. 40 of 812 subjects reported treatment-related adverse experiences.

Local Adverse Reactions Occurring at an Incidence \geq 1% in Clinical Trials

| | |
|---------------------|--|
| Ointment (n=812) | Burning (1.6%) Pruritus (1%) Skin Atrophy (1%) |
|---------------------|--|

Less Common Local Adverse Reactions in Clinical Trials (<1%)

Infections and infestations: furuncle

Nervous system disorders: paraesthesia

Skin and subcutaneous tissue disorders: pain of skin

The following local adverse reactions have been reported infrequently when other topical dermatologic corticosteroids have been used as recommended. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, miliaria.

Hypothalamic-pituitary-adrenal (HPA) axis suppression has been reported with topical corticosteroids. Manifestations of HPA axis suppression include increased weight / obesity, delayed weight gain / growth retardation in children, Cushing's syndrome, cushingoid features (e.g., moon face, central obesity), HPA disorder, decreased endogenous cortisol levels, hyperglycemia / glucosuria, hypertension, osteoporosis, cataract, glaucoma, steroid withdrawal syndrome (see WARNINGS AND PRECAUTIONS).

Systemic adverse reactions, such as vision blurred, have also been reported with the use of topical corticosteroids.

Posterior subcapsular cataracts have been reported following systemic use of corticosteroids.

DRUG INTERACTIONS

Drug-Drug Interactions

Interactions with other drugs have not been established.

Co-administered drugs that can inhibit CYP3A4 (e.g., ritonavir, itraconazole) have been shown to inhibit the metabolism of corticosteroids leading to increased systemic exposure. The extent to which this interaction is clinically relevant depends on the dose and route of administration of the corticosteroids and the potency of the CYP3A4 inhibitor.

Drug-Food Interactions

Interaction with food has not been established.

Drug-Herb Interactions

Interaction with herbs has not been established.

Drug-Laboratory Interactions

Interaction with laboratory testing has not been established.

Drug-Lifestyle Interactions

Interaction with lifestyle has not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

- Patients/caregivers should be instructed to use TEVA-MOMETASONE for the minimum amount of time necessary to achieve the desired results because of the potential for corticosteroids to suppress the hypothalamic-pituitary-adrenal (HPA) axis and cause skin atrophy (See WARNINGS AND PRECAUTIONS).
- TEVA-MOMETASONE is for topical use only and not for use on mucous membranes.
- Use in pediatric patients is not recommended. Pediatric patients may be more susceptible to local and systemic toxicity from equivalent doses because of their larger skin surface to body weight ratios.
- Geriatric patients may be more susceptible to the potential effects of systemic absorption. Decreased hepatic or renal function in the elderly may delay elimination.

Recommended Dose and Dosage Adjustment

Apply a thin film of TEVA-MOMETASONE to the affected skin areas once daily.

TEVA-MOMETASONE should be used **on the face, axillae or scrotum for a maximum of 5 days**.

TEVA-MOMETASONE should be used on the **body for a maximum of 3 weeks**. If no improvement is seen within 2 weeks, reassessment of diagnosis and treatment may be necessary.

Therapy should be discontinued when control is achieved. Continue an emollient as maintenance therapy.

Pediatrics (< 18 years of age): Safety and efficacy of mometasone have not been established in pediatric patients. (see WARNINGS AND PRECAUTIONS – Special Populations, Pediatrics).

Geriatrics (≥ 65 years of age): Mometasone should be used with caution due to increased risk of renal or hepatic impairment in this population. The minimum quantity should be used for the shortest duration to achieve the desired clinical benefit (see WARNINGS AND PRECAUTIONS– Special Populations, Geriatrics).

Patients with renal/hepatic impairment: The minimum quantity should be used for the shortest duration to achieve the desired clinical benefit (see WARNINGS AND PRECAUTIONS-Special Populations, Patients with renal/hepatic impairment).

Missed Dose

Any missed dose should be applied as soon as possible after the missed dose is remembered. If this is close to the scheduled application time or the next dose, the subject should wait and apply the next scheduled dose. The usual schedule should be resumed thereafter.

Administration

Do not use occlusive dressings.

TEVA-MOMETASONE should not be applied to mucous membranes.

OVERDOSAGE

| |
|---|
| For management of a suspected drug overdose, contact your regional Poison Control Centre. |
|---|

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects. Excessive prolonged use or misuses may suppress hypothalamic-pituitary-adrenal (HPA) axis function, resulting in secondary adrenal insufficiency. If symptoms of HPA axis suppression occur, TEVA-MOMETASONE should be gradually discontinued by reducing the frequency of application or by substituting a less potent corticosteroid, as clinically indicated. If toxic effects occur, TEVA-MOMETASONE should be discontinued and symptomatic therapy administered (see WARNINGS AND PRECAUTIONS).

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

TEVA-MOMETASONE (mometasone furoate) has anti-inflammatory, antipruritic and vasoconstrictive actions. The exact mechanism, however, of corticosteroids in each disease is uncertain. Mometasone furoate, has been shown to have topical (dermatologic) and systemic pharmacologic and metabolic effects characteristic of this class of drugs.

Pharmacodynamics

In a study of the effects of mometasone furoate ointment on the HPA axis, 15 g were applied BID for seven days to patients with psoriasis or atopic dermatitis. The ointment was applied without occlusion to at least 30% of body surface. The results suggest that the drug caused a slight reduction of urinary-free cortisol. However, this change was not considered clinically important since it was not accompanied by subnormal levels of plasma cortisol or 17-OHC.

Pharmacokinetics

A percutaneous absorption study with radio-labeled ³H-mometasone furoate ointment was conducted in adult male volunteers with intact skin. Based on the amounts of radioactivity excreted after an eight-hour application of the active ointment and analysis of urine and feces, approximately 0.7% of the applied dose was absorbed systemically without occlusion.

STORAGE AND STABILITY

Store between 15° and 30°C.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Each gram TEVA-MOMETASONE Ointment 0.1% contains 1 mg mometasone furoate. Non medicinal ingredients include: hexylene glycol, propylene glycol monostearate, white petrolatum and white wax.

TEVA-MOMETASONE Ointment 0.1% is supplied in 15 g and 50 g tubes.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: mometasone furoate

Chemical name: 9a,21-Dichloro-11b,17-dihydroxy-16a-methylpregna-1,4-diene-3,20-dione 17-(2-furoate)

Molecular formula and
molecular mass: $C_{27}H_{30}Cl_2O_6$ and a
molecular weight of 521.4 g/mol

Structural formula:



Physicochemical properties: Mometasone furoate is a white to off-white powder practically insoluble in water, slightly soluble in octanol and moderately soluble in ethyl alcohol.

CLINICAL TRIALS

Efficacy Studies: Mometasone Furoate Ointment 0.1%

Psoriasis - In two bilateral-paired comparison trials, the efficacy of BID applications of mometasone furoate ointment in concentrations of 0.1% and 0.05% was compared to that of betamethasone valerate ointment also applied BID for 14 days. Results showed that the 0.1% formulation of mometasone furoate ointment was significantly ($P<0.05$) more effective than betamethasone valerate ointment^a. As demonstrated by the physician's global evaluation of change in disease status, 60% of patients responded more favorably to mometasone furoate ointment 0.1%, while 13% experienced a comparable response in the betamethasone valerate-treated group. Improvement from baseline in total disease sign score was 51% and 40% for mometasone furoate ointment 0.1% and betamethasone valerate ointment, respectively. Furthermore, these results also demonstrated that mometasone furoate ointment 0.05% was superior to betamethasone valerate ointment but not as effective as the 0.1% mometasone furoate ointment formulation.

In a third bilateral-paired comparative study of mometasone furoate ointment 0.1% and betamethasone dipropionate ointment^b applied BID for 14 days, percent improvement in total disease scores was similar between the two preparations, 63% and 58% for mometasone furoate ointment 0.1% and betamethasone dipropionate ointment, respectively. However, 38% of patients responded more favorably to mometasone furoate ointment 0.1% while 3% responded better to betamethasone dipropionate ointment.

Furthermore, three randomized, multicentric, parallel group studies were conducted in patients with psoriasis to compare the efficacy of mometasone furoate ointment 0.1% applied QD to that of triamcinolone acetonide^c applied BID, fluocinolone acetonide^d applied TID or to that of the vehicle alone applied QD for 21 days. Mometasone furoate ointment 0.1% was significantly ($P<0.01$) better than triamcinolone acetonide, fluocinolone acetonide and the vehicle as demonstrated by the percent improvement in total disease sign scores. The superior efficacy of mometasone furoate ointment applied QD was observed despite the more frequent administrations of the two comparative agents. Physician's global evaluation of disease status at endpoint analysis also confirmed that mometasone furoate ointment 0.1% was significantly ($P<0.01$) more effective than triamcinolone acetonide, fluocinolone acetonide or the vehicle alone in the treatment of patients with psoriasis.

a VALISONE® Ointment

b DIPROSONE® Ointment

c KENALOG® ER Squibb & Sons, Inc., Princeton, NJ USA

d SYNALAR® Syntax Laboratories, Palo Alto, CA USA

e BETNOVATE® Ointment, Glaxo Laboratories, UK

Two additional studies in psoriatic patients compared QD applications of mometasone furoate ointment 0.1% with QD applications of betamethasone dipropionate 0.05% and BID applications of betamethasone valerate 0.1%^e respectively for three weeks. Mometasone furoate ointment 0.1% QD was significantly ($P<0.01$) more effective than betamethasone valerate BID and comparable to betamethasone dipropionate QD as demonstrated by percent improvement in total disease sign scores at endpoint analysis. Physician's overall evaluation of disease status also indicated that mometasone furoate ointment was significantly ($P<0.01$) more effective than betamethasone valerate in the treatment of psoriasis. At the end of the three-week study period, mean scores were indicative of marked to moderate improvement in most patients treated with mometasone furoate ointment. Comparable improvement was effected with betamethasone dipropionate and moderate to slight improvement was observed in the betamethasone valerate treated group.

Atopic Dermatitis -- Patients with atopic dermatitis participated in a bilateral-paired comparative study, which evaluated the efficacy of mometasone furoate ointment 0.1% against that of betamethasone valerate ointment. Results demonstrated that mometasone furoate ointment 0.1% was equivalent in activity to betamethasone valerate ointment when both agents were applied BID. Another three randomized, multicentric parallel-group studies compared the efficacy of mometasone furoate ointment 0.1% QD with that of betamethasone valerate ointment BID, the ointment vehicle alone applied QD, or hydrocortisone butyrate ointment 0.1%^f applied BID for three weeks. In these studies, mometasone furoate was equivalent to the known standard agents, betamethasone valerate and hydrocortisone butyrate, even though mometasone furoate was applied less frequently than each of these comparatives. Percent improvement in total disease sign score at end-point analysis in the three studies were 82%, 83% and 60%, respectively, for mometasone furoate ointment 0.1% as compared to 79%, 24% and 46% for betamethasone valerate ointment, the vehicle and hydrocortisone butyrate, respectively ($P<0.01$). Furthermore, global scores at endpoint reflected marked improvement in the mometasone furoate and betamethasone valerate-treatment groups, moderate improvement in the hydrocortisone-treated group and slight improvement in the vehicle-treated group.

Corticosteroid-Responsive Dermatoses -- In three parallel-group studies, the efficacy of mometasone furoate ointment 0.1% was compared to that of betamethasone valerate 0.05% and clobetasone butyrate 0.025% in the treatment of various corticosteroid-responsive dermatoses. Mometasone furoate ointment was applied QD while the comparative agents were each applied BID for three weeks. After one treatment week, improvement in disease signs ranged from 58% to 90% with QD mometasone furoate administration, 52% to 77% with BID application of betamethasone valerate and 69% with BID administration of clobetasone butyrate. By treatment end, percent improvement averaged 93% for mometasone furoate, 89% and 93% for betamethasone

valerate and 90% for clobetasone butyrate. At endpoint evaluation, global scores indicated disease clearance in the majority of mometasone-treated patients; marked improvement was observed in most patients treated with betamethasone valerate or clobetasone butyrate.

Onset of Action: Ointment -- Mometasone furoate ointment 0.1% QD also had a rapid onset of action in psoriatic patients as evidenced by percent improvement from baseline in total disease sign/symptom scores after one treatment week (ranging from 38% to 59%). Percent improvements for comparative agents were triamcinolone acetonide (28%), fluocinolone acetonide (33%), betamethasone dipropionate (23%), betamethasone valerate (56%) and vehicle alone (43%). In two of these studies mometasone furoate was significantly more effective than triamcinolone acetonide or fluocinolone acetonide at Day 4 evaluation ($P < 0.01$).

The effects of mometasone furoate ointment 0.1% in the treatment of patients with atopic dermatitis also were rapid in onset as demonstrated by mean percent improvement and mean global evaluation scores at Day 4 and Week 1. Mometasone furoate-treated patients showed an improvement in total sign/symptom score that ranged from 27% to 47% at Day 4 and 51% to 64% at Week 1. In comparison, hydrocortisone butyrate and betamethasone valerate demonstrated 17% and 43% improvement, respectively, at Day 4 and 24% and 65%, respectively, at Week 1. Global scores at one-week indicated moderate improvement in patients treated with mometasone furoate or betamethasone valerate and slight improvement in those treated with hydrocortisone butyrate.

Safety Studies - In a study of the effects of mometasone furoate ointment on the HPA axis, 15 g were applied BID for seven days to patients with psoriasis or atopic dermatitis. The ointment was applied without occlusion to at least 30% of body surface. The results suggest that the drug caused a slight reduction of urinary-free cortisol. However, this change was not considered clinically important since it was not accompanied by subnormal levels of plasma cortisol or 17-OHC.

The results of other local and systemic safety studies also showed that mometasone furoate ointment 0.1% has minimal percutaneous absorption and do not cause adrenal suppression. In other investigations, mometasone furoate ointment 0.1% demonstrated minimal potential for irritation, sensitization, photocontact allergenicity and phototoxic reactions when used as recommended. Furthermore, when compared to hydrocortisone ointment 0.1% mometasone furoate ointment 0.1% has a low atrophogenic potential. No clinical meaningful changes in laboratory test values were observed with mometasone furoate ointment.

Comparative Bioavailability Study

In a comparative bioequivalence study between **TEVA*-MOMETASONE 0.1%** Ointment and the Canadian reference product, the relative potency of each product was measured using a ChromaMeter. This device measures the degree of local vasoconstrictor response by measuring the degree of skin blanching produced by each drug.

The following table reveals the results of the skin blanching measurements. Both products demonstrated comparable vasoconstrictor (therapeutic) activity.

**Mean ChromaMeter Results comparing
TEVA*-MOMETASONE 0.1% Ointment (Test)
with the Canadian Reference Product (Reference)**

| Evaluation | N | Test | Reference: | Ratio | 90% Confidence Interval ² | |
|---|----|--|---|------------------|--------------------------------------|--------------|
| | | TEVA*- MOMETASONE 0.1% Ointment (Teva Canada Limited) | Elocom 0.1% Ointment (Schering Canada) | (%) ¹ | Lower (%) | Upper (%) |
| ChromaMeter Reading (Skin blanching) | 38 | 13 | 13.3 | 98 | 88 | 108.5 |

1: Ratio percent calculated as Test/Reference x 100

2: Confidence interval on the ratio

DETAILED PHARMACOLOGY

Pre-Clinical Data

Pharmacodynamics -- The pharmacologic profile of mometasone furoate was determined by standard laboratory methods. Relative to betamethasone valerate, the anti-inflammatory activity and anti-psoriatic activity of mometasone furoate were evaluated in mice and guinea pigs, respectively. Hypothalamic-pituitary-adrenal (HPA) axis suppression, thymolysis and skin atrophy were evaluated in mice.

In the croton oil assay in mice, mometasone furoate ($ED_{50} = 0.02 \mu\text{g}/\text{ear}$) was equipotent to betamethasone valerate after single application, and was approximately eight times as potent as betamethasone valerate after five daily applications (ED_{50} values = $0.002 \mu\text{g}/\text{ear}/\text{day}$ vs $0.014 \mu\text{g}/\text{ear}/\text{day}$). In guinea pigs, mometasone furoate was approximately twice as potent as betamethasone valerate in reducing M. Ovalis-induced epidermal acanthosis after 14 daily applications.

With respect to pharmacologic activities commonly associated with corticosteroids, mometasone furoate ($ED_{50} = 5.3 \mu\text{g}/\text{ear}/\text{day}$) was less potent than betamethasone valerate ($ED_{50} = 3.1 \mu\text{g} /\text{ear}/\text{day}$) in suppressing the HPA axis in mice after five daily applications. In the thymolysis assay, mometasone furoate ($ED_{50} = 26.6 \mu\text{g} /\text{ear}/\text{day}$) was approximately two times as potent as betamethasone valerate ($ED_{50} = 51.6 \mu\text{g} /\text{ear}/\text{day}$) when applied topically, and following subcutaneous administration for five days, mometasone furoate ($ED_{50} = 11.2 \mu\text{g} /\text{mouse}$) was approximately six times as potent as betamethasone valerate ($ED_{50} = 59.8 \mu\text{g} /\text{mouse}$). At doses five to 5000 times the effective anti-inflammatory doses, mometasone furoate was three to eight times more potent than betamethasone valerate with respect to skin thinning in mice. Based on the ratio of systemic potency (HPA suppression or thymolysis) to topical anti-inflammatory potency, the therapeutic indexes for mometasone furoate were approximately three to ten times greater than those for the comparative, betamethasone valerate. Therefore, mometasone furoate would be expected to have a superior safety margin to that of betamethasone valerate.

Pharmacokinetics -The percutaneous absorption and excretion of ^3H mometasone furoate ointment was evaluated in rats, rabbits and dogs with doses ranging from 5.2 to $22 \mu\text{g}/\text{cm}^2$. Additionally, the tissue distribution of absorbed radioactivity was determined in rabbits.

Systemic absorption of ^3H -mometasone furoate was minimal in all species studied, ranging from approximately 2% in dogs to 6% in rabbits over a 5 to 7-day period. The cream and ointment formulations were comparable with respect to systemic absorption. Plasma levels were low ranging from <0.1 to $<1 \text{ ng}/\text{ml}$. Less than 1.3% of the applied dose was excreted in urine of all species and from 1.5 to 4.2% was excreted in feces. Characterization of urinary metabolites was not possible due to the low levels of drug in urine. However, it is well known that corticosteroids are metabolized to inactive water-soluble substances such as sulfate esters or glucuronides and are

excreted as such. In rabbits, there was no unusual accumulation of radioactivity in any tissue.

TOXICOLOGY

Toxicology -- A program consisting of evaluation of local and systemic toxicity, reproductive toxicity, genetic toxicity, dermal irritation and sensitization potential and ocular irritation was conducted to determine the safety of mometasone furoate cream and ointment. Acute toxicity was evaluated in mice, rats and dogs including young (21-day old) mice and rats. Repeated dose toxicity was evaluated in rats, rabbits and dogs by subcutaneous and/or topical routes. Reproduction studies were conducted in rats and rabbits and included evaluation of teratology, peri and post- natal development and general reproductive performance. Sensitization potential was determined in guinea pigs and dermal and ocular irritation were evaluated in rabbits. In vitro and in vivo genetic toxicology studies were conducted to evaluate potential mutagenicity and clastogenicity (capacity to induce chromosomal changes).

The acute subcutaneous LD₅₀ values of mometasone furoate were determined to be between 200 and 2000 mg/kg in mice, 2000 mg/kg or greater in rats and >200 mg/kg in dogs. Following oral administration the LD₅₀ values were >2000 mg/kg in mice and rats. As expected, the LD₅₀ values for young (21-day old) mice and rats were 2 to 20 times lower than those for adult animals.

Following repeated administration of mometasone furoate in rats, rabbits and dogs at doses up to 670 times the anticipated maximum human dose for up to 6 months, findings were typical of corticosteroid administration in all species. These included (1) slight reduction in body weight gain, (2) skeletal muscle wasting, (3) abdominal distention, (4) decrease in lymphocytes and eosinophils and increase in neutrophils, (5) increase in serum transaminases (ALT and AST), cholesterol and triglycerides, (6) lipemia, and (7) organ changes (atrophy of spleen and thymus, local skin thinning, increased liver and kidney weights and reduced osteogenesis). These changes were generally observed more frequently or more severe in animals receiving the comparative agent, betamethasone valerate. No unusual systemic effects were observed with either drug. Dermal responses to repeated application of mometasone furoate or betamethasone valerate cream were limited to transient episodes of slight to moderate erythema, skin wrinkling, desquamation and the presence of papules and/or pustules.

In reproduction studies, mometasone furoate produced effects which are known to be associated with corticosteroids and/or progestational agents such as reduced maternal body weight gain, suppression of fetal growth, delayed ossification, umbilical hernias, prolonged gestation, difficult and prolonged labor and inability to deliver.

In genetic toxicity studies, mometasone furoate was not mutagenic in bacteria (Ames

test) or mammalian (mouse lymphoma) cells and was not clastogenic in the mouse micronucleus test.

REFERENCES

1. Schering Corporation, Kenilworth, N.J. Percutaneous Absorption of ^3H -SCH 32088 (Ointment) in the Rabbit, (P-5039); 1984.
2. Schering Corporation, Kenilworth, N.J. Percutaneous Absorption and Tissue Distribution of ^3H -SCH 32088 (Ointment) in the Rabbit, (P-5027); 1984.
3. Schering Corporation, Kenilworth, N.J. Percutaneous Absorption of ^3H -SCH 32088 in the Rabbit, (P-5133); 1986.
4. Schering Corporation, Kenilworth, N.J. Percutaneous Absorption of ^3H -SCH 32088 in the Dog, (P-5045); 1985.
5. Schering Corporation, Kenilworth, N.J. Acute Oral and Subcutaneous Toxicity Studies of SCH 32088 in Rats and Mice, (P-4865); 1982.
6. Schering Corporation, Kenilworth, N.J. Acute Subcutaneous Toxicity Study of SCH 32088 in Young (21-Day Old) Mice, (P-5165); 1986.
7. Schering Corporation, Kenilworth, N.J. Acute Subcutaneous Toxicity Study of SCH 32088 in Young (21-Day Old) Rats, (P-5168); 1986.
8. Schering Corporation, Kenilworth, N.J. Acute Subcutaneous Toxicity Study of SCH 32088 in Dogs, (P-4868); 1982.
9. Schering Corporation, Kenilworth, N.J. Two-Week Subcutaneous Toxicity Study of SCH 32088 in Rats, (P-4895); 1983.
10. Schering Corporation, Kenilworth, N.J. Three-Week Dermal Toxicity Study of SCH 32088 Ointment in Rabbits with Intact or Abraded Skin, (P-4919); 1983,.
11. Schering Corporation, Kenilworth, N.J. Three-Month Dermal Toxicity Study of SCH 32088 Ointment (0.1%) in Rabbits with Intact Skin, (P-5010); 1985.
12. Schering Corporation, Kenilworth, N.J. Two-Week Subcutaneous Toxicity Study of SCH 32088 in Beagle Dogs, (P-4883); 1983.
13. Schering Corporation, Kenilworth, N.J. Six-Month Dermal Toxicity Study of SCH 32088 Ointment (0.1%) in Dogs with Intact Skin, (P-5013); 1985.
14. Schering Corporation, Kenilworth, N.J. Dermal Teratology (Segment II) Study of

- SCH 32088 Ointment (0.1%) in Rats, (P-5054); 1985.
15. Schering Corporation, Kenilworth, N.J. Dermal Teratology (Segment II) Study of SCH 32088 Ointment (0.1%) in Rats, (P-5066); 1985.
 16. Schering Corporation, Kenilworth, N.J. Perinatal and Postnatal Reproduction (Segment III) Study of SCH 32088 in Rats, (P-5164); 1986.
 17. Schering Corporation, Kenilworth, N.J. Fertility and General Reproduction Study (Segment I) of SCH 32088 in Rats, (P-5174); 1987.
 18. Schering Corporation, Kenilworth, N.J. Dermal Sensitization Study of SCH 32088 in Guinea Pigs, (P-4879); 1983.
 19. Schering Corporation, Kenilworth, N.J. Acute Ocular Irritation Study of SCH 32088 Ointment in Rabbits, (P-4933); 1983.
 20. Schering Corporation, Kenilworth, N.J. Acute Ocular Irritation Study of SCH 32088 Cream in Rabbits, (P-4950); 1984.
 21. Schering Corporation, Kenilworth, N.J. Salmonella/ Mammalian Microsome Bioassay of SCH 32088. (P-4988); 1984.
 22. Schering Corporation, Kenilworth, N.J. L5178 TK +/-, TK -/- Mouse Lymphoma Cell Mutagenicity Assay of SCH 32088. (P-5011); 1984.
 23. Schering Corporation, Kenilworth, N.J. Mouse Bone Marrow Erythrocyte Micronucleus Assay of SCH 32088. (P-5050); 1985.
 24. Nagabhushan, N. et al, Percutaneous Absorption of ³H-SCH 32088 in Male Volunteers, (C84-103), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
 25. Medansky, R., Bilateral Paired Comparison Study of SCH 32088 Ointment 0.1% and VALISONE Ointment 0.1% in Psoriasis, (C83066), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
 26. Medansky, R., Bilateral Paired Comparison Study of SCH 32088 Ointment 0.05% and VALISONE Ointment 0.1% in Psoriasis, (C83-067), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
 27. Medansky, R., Bilateral Paired Comparison Study of SCH 32088 Ointment 0.1% and DIPROSONE Ointment 0.05% in Psoriasis, (C84-03), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.

28. Liebsohn, E., et al, Single-blind Cooperative Efficacy and Safety Study of SCH 32088 Ointment 0.1% QD and KENALOG Ointment 0.1% BID in Psoriasis, (C84-043), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
29. Lasser, A., et al, Single-blind Cooperative Efficacy and Safety Study of SCH 32088 Ointment 0.1% QD and SYNALAR Ointment 0.025% TID in Psoriasis, (C84-047), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
30. Kanof, N., et al, Double-blind Cooperative Efficacy and Safety Study of SCH 32088 Ointment 0.1% QD and Its Vehicle QD in Psoriasis, (C84-055), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
31. Daniel, J. and Thivolet, J., Single-blind Efficacy and Safety Study in Psoriasis Patients Comparing Once Daily Applications of Mometasone Furoate Ointment 0.1 and DIPROSONE Ointment 0.05%, (I86-211-01, 02), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
32. Rosenthal, D. and Duke, E., Single-blind Efficacy and Safety Study in Psoriasis Patients Comparing Once Daily Applications of Mometasone Furoate Ointment 0.1% and Twice Daily Applications of BETNOVATE® Ointment 0.1%, (I86-308-01, 02), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
33. Hanifin, J., Bilateral Paired Comparison Study of SCH 32088 Ointment 0.1% and VALISONE Ointment 0.1% in Atopic Dermatitis, (C84-020), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
34. Roth, H., et al, Single-blind Cooperative Efficacy and Safety Study of SCH 32088 Ointment 0.1% QD and VALISONE Ointment 0.1% BID in Atopic Dermatitis, (C84- 048), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
35. Rex, I., et al, Double-blind Cooperative Efficacy and Safety Study of SCH 32088 Ointment in Psoriasis Comparing 0.1% and Its Vehicle QD in Atopic Dermatitis, (C84- 065), Mometasone Furoate Ointment 0.1%, Health Registration Dossier, Schering International; 1987.
36. Cerio, R. and MacDonald, D.M., Single-blind Efficacy and Safety Study in Atopic Dermatitis Patients Comparing Once Daily Applications of Mometasone Furoate Ointment 0.1% and Twice Daily Applications of LOCOID® Ointment 0.1%, (I86- 309- 01,02), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.

37. Jaimovich, L., Single-blind Efficacy and Safety Study in Patients with Steroid-responsive Dermatoses Comparing Once Daily Applications of Mometasone Furoate Ointment 0.1% and Twice Daily Applications of BETNOVATE® Ointment 0.1%, (I86-118), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
38. Moncada, B., Single-blind Efficacy and Safety Study in Patients with Steroid-responsive Dermatoses Comparing Once Daily Applications of Mometasone Furoate Ointment 0.1% and Twice Daily Applications of EUMOVATE® Ointment 0.05%, (I86-119), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
39. Meinicke, K. et al, Single-blind Efficacy and Safety Study in Patients with Steroid-responsive Dermatoses Comparing Once Daily Applications of Mometasone Furoate Ointment 0.1% and Twice Daily Applications of Betnovate® Ointment 0.1%, (I87-211- 01, 02, 03, 04), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
40. Katz, H., Systemic Tolerance to Topical Applications of SCH 32088 Ointment 0.1% in In-patients with Psoriasis or Atopic Dermatitis, (C84-052), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
41. Kaidbey, K., Evaluation of Photocontact Allergenicity Potential of SCH 32088 Ointment 0.1% and Its Vehicle, (C84-039), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
42. Kaidbey, K., Evaluation of Phototoxicity Potential of SCH 32088 Ointment 0.1% and its Vehicle, (C84-040), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
43. Jordan, W., A Cumulative Irritancy Assay To Determine the Relative Irritation Potential of SCH 32088 Ointment, (C83-091) Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
44. Jordan, W., Evaluation of the Irritation and Sensitization Potentials of SCH 32088 Ointment and Its Vehicle, (C83-092), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1987.
45. Bressinck, R., et al, Effect of Mometasone Furoate Ointment 0.1% and Hydrocortisone Ointment 1% on Adrenocortical Function in Psoriasis Patients, (C85-077), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
46. Katz, H., Bilateral-paired Comparison Study of the Atrophogenic Potential of Mometasone (SCH 32088) Ointment, 0.1% and

- Hydrocortisone Ointment, 1.0% (HYTONE® in Psoriasis, (C86-064-01), Mometasone Furoate Ointment 0.1% Health Registration Dossier, Schering International; 1988.
47. Belsito DV, Baer RL, Schultz JM, Thorbecke GJ. Relative lack of systemic effects of mometasone furoate on Langerhans Cells of mice after topical administration as compared with other glucocorticosteroids. *J Invest Dermatol* 1988 91:219-23.
 48. Bressinck R, Williams J, Peets E. Comparison of the effect of mometasone furoate ointment 0.1%, and hydrocortisone ointment 1%, on adrenocortical function in psoriasis patients. *Today's Therapeutic Trends* 5:25-34, 1988.
 49. Medansky RS, Brody NI, Kanof NB, Russo GJ, Peets EA. Clinical investigations of mometasone furoate - a novel nonfluorinated, topical corticosteroid. *Topical Corticosteroid Seminars in Dermatology*. 1987 6(2):94-100.
 50. Medansky RS, Bressinck R, Cole GW, Deeken JH, Ellis CN, Guin JD, et al. Mometasone furoate ointment and cream 0.1 percent in treatment of psoriasis: comparison with ointment and cream formulations of fluocinolone acetonide 0.025 percent and triamcinolone acetonide 0.1 percent. 1988 *Cutis* 41:480-5.
 51. Rosenthal D, Duke E. A clinical investigation of the efficacy and safety of mometasone furoate ointment 0.1% vs betamethasone valerate ointment 0.1% in the treatment of psoriasis. 1988 *Curr Ther Res* 44(5):790-801.
 52. Product Monograph Elocom, Merck Canada Inc., December 21, 2017 (control# 210612)

PART III: CONSUMER INFORMATION

**PrTEVA-MOMETASONE
Mometasone Furoate Ointment, 0.1%**

This leaflet is part III of a three-part "Product Monograph" published when TEVA-MOMETASONE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about TEVA-MOMETASONE. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

In adult, TEVA-MOMETASONE:

- is used for the relief of swelling and itching caused by skin conditions like psoriasis and atopic dermatitis (a type of eczema).
- can be used for a maximum of:
 - 5 days on the face, scalp, skin-fold areas and groin
 - 3 weeks on the body

What it does:

TEVA-MOMETASONE, has anti-inflammatory and vasoconstrictive actions (makes blood vessels constrict) to help relieve swelling and itching. The exact mechanism of action is not known.

When it should not be used:

Do not use TEVA-MOMETASONE, if you:

- are allergic to mometasone furoate, other corticosteroids or to any of the ingredients of TEVA-MOMETASONE. (*See section, What the important nonmedicinal ingredients are*).
- have bacterial, fungal, parasitic, viral skin infection (e.g. herpes simplex, chickenpox), tuberculous or syphilis skin lesions, or skin reaction following a recent vaccination
- have acne
- have rosacea (a facial skin condition where the nose, cheeks, chin, forehead or entire face are unusually red, with or without tiny visible blood vessels, bumps (papules) or pus-filled bumps (pustules)
- have itchy skin which is not inflamed

Do not apply TEVA-MOMETASONE in the eyes.

What the medicinal ingredient is:

Mometasone furoate

What the important nonmedicinal ingredients are:

- hexylene glycol

- propylene glycol monostearate
- white petrolatum
- white wax

What dosage forms it comes in:

- TEVA-MOMETASONE Ointment 0.1% is supplied in 15 g and 50 g tubes

WARNINGS AND PRECAUTIONS

Apply just enough TEVA-MOMETASONE to cover the areas. TEVA-MOMETASONE can get into the blood and cause side effects.

Always follow your doctor's instructions.

TEVA-MOMETASONE is NOT recommended for use with airtight bandage.

TEVA-MOMETASONE is more likely to cause side effects when used:

- over large areas
- on sensitive areas such as the face, scalp, skin fold areas like the armpit and groin
- on broken skin
- for a long time

Inform any doctor you consult that you are using or have previously used corticosteroids.

BEFORE using TEVA-MOMETASONE, talk to your doctor or pharmacist if:

- you have any skin disease around a leg ulcer; use of a topical corticosteroid may increase the risk of an allergic reaction or an infection around the ulcer.
- you are currently treating an infection using an antifungal or antibacterial agent.
- you have other inflammatory skin diseases in the leg as a result of impaired circulation (such as stasis dermatitis).
- you are pregnant or planning to become pregnant.
- you are breastfeeding. It is not known if TEVA-MOMETASONE will appear in breast milk. You should only use TEVA-MOMETASONE while breastfeeding if you and your doctor decide if the benefits to the mother outweigh the risks to the baby. If you do breastfeed when using TEVA-MOMETASONE, do not apply it on your breasts to ensure the infant does not accidentally get it in their mouth.
- you have problems with your kidney and liver. You may need to use a smaller amount of TEVA-MOMETASONE or use it less often.

While using TEVA-MOMETASONE, talk to your doctor

or pharmacist if:

- you develop any skin infection
- you have an allergic reaction
- you develop significant skin irritation
- you experience skin thinning or softening
- your condition worsens or does not improve

Care should be taken when applying TEVA-MOMETASONE to the face or in skin-fold areas, such as the groin and the armpit since these areas are more prone to skin thinning.

Use with caution when applying TEVA-MOMETASONE near the eyes.

Children absorb larger amounts of topical corticosteroids and therefore, may be more likely to develop side effects. **TEVA-MOMETASONE is not recommended for use in children under 18 years of age.**

Do NOT cover the affected skin or scalp areas with airtight bandages.

INTERACTIONS WITH THIS MEDICATION

It is NOT known whether TEVA-MOMETASONE interacts with other medication. Some medicines may affect how TEVA-MOMETASONE works and may make it more likely that you will have side effects. Some of these medicines may include:

- Ritonavir (for HIV).
- Itraconazole (for fungal infections).

Tell your doctor or pharmacist about all your other medications, including medicines that you bought without a prescription and natural health products.

PROPER USE OF THIS MEDICATION

Use the minimum quantity of TEVA-MOMETASONE for the shortest amount of time necessary to achieve the desired results. This is especially important if you are 65 years or older or have liver or kidney disease.

TEVA-MOMETASONE is for use on the skin and scalp only. It is **NOT** for use in the eyes or other mucous membranes.

Usual Adult dose:

TEVA-MOMETASONE Ointment: Apply a thin film to the affected skin areas once daily or as directed by your doctor.

Use for a maximum of:

- 5 days on the face, skin-fold areas and groin.
- 3 weeks on the body. If your condition worsens or no improvement is seen within 2 weeks, contact your doctor.

Use TEVA-MOMETASONE only as directed by your health care provider. **Do NOT use more of it, do NOT use it more often and do NOT use it for a longer period of time than your health care provider recommended.** Using too much TEVA-MOMETASONE may increase your chances of unwanted and sometimes dangerous side effects.

This medication has been prescribed specifically for you. Do NOT give it to anyone else. It may harm them, even

if their symptoms seem to be similar to yours.

Overdose:

If you think you have taken too much TEVA-MOMETASONE, contact your health care practitioner, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed dose:

If you forget to use TEVA-MOMETASONE, apply it as soon as you remember. If it is close to the time scheduled to apply your next dose, wait and apply your next scheduled dose and then continue as before. Do not apply extra TEVA-MOMETASONE to make up for missed doses.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- your condition worsens or does not improve
- skin sensations such as:
 - burning, tingling, or stinging
 - strong itching sensation
 - softening and thinning of the skin
 - inflammation of hair follicles
 - acne-like reactions

Very rarely the following may occur:

- aggravation of the disease
- dry skin, abnormal redness, appearance of boils, aggravated allergy, dermatitis (swelling of the skin), increased lesion size
- nausea

Additionally, the following side effects have been found to occur with the use of other topical corticosteroids:

- infection or signs of infection, irritation
- unwanted hair, lightening of skin color
- dermatitis (swelling of the skin) near or around the mouth, allergic contact dermatitis
- stretch marks
- heat rash

TEVA-MOMETASONE can cause abnormal blood and urine test results. Your doctor will decide when to perform tests and will interpret the results.

| SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM | | | | |
|--|--|-------------------------------------|--------------|---|
| Symptom / effect | | Talk with your doctor or pharmacist | | Stop taking drug and get immediate medical help |
| | | Only if severe | In all cases | |
| Un-common | Allergic reaction: chills, fever, muscle aches or pains or other flu-like symptoms occurring with or before a skin rash | | | ✓ |
| | Cushing's Syndrome: weight gain, moon face / rounding of the face and obesity | | | ✓ |
| Unknown | Steroid Withdrawal Syndrome: weight loss, fatigue, nausea, diarrhea and abdominal pain | | ✓ | |
| | Hyperglycemia (increased blood sugar): frequent urination, thirst and hunger | | ✓ | |
| | Glucosuria (sugar in urine): excessive or sweet smelling urine | | ✓ | |
| | Hypertension (high blood pressure): headaches, vision disorders, nausea and vomiting | | ✓ | |
| | Osteoporosis: weakening of the bones potentially leading to an increased risk of bone fracture | | ✓ | |

| | | | |
|---|--|---|--|
| Glaucoma, Cataracts, or central serous chorioretinopathy (CSCR): blurred vision, increased pressure in your eyes, eye pain, distorted vision | | ✓ | |
|---|--|---|--|

*This is not a complete list of side effects. For any unexpected effects while taking **TEVA-MOMETASONE ointment** contact your doctor or pharmacist.*

HOW TO STORE IT

Store between 15° and 30°C

Do **NOT** use if past expiry date on the label.

Keep out of reach and sight of children.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

If you want more information about TEVA-MOMETASONE:

- Talk to your healthcare professional
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website <http://www.tevacanada.com>; or by calling 1-800-268-4127 ext. 3; or email druginfo@tevacanada.com.

This leaflet was prepared by:
Teva Canada Limited
30 Novopharm Court
Toronto, Ontario
M1B 2K9
Canada

www.tevacanada.com

Last revised: May 7, 2018

THIS IS A REPRESENTATION OF AN ELECTRONIC RECORD THAT WAS SIGNED ELECTRONICALLY AND THIS PAGE IS THE MANIFESTATION OF THE ELECTRONIC SIGNATURE

Teva Canada Limited

Teva-Mometasone Ointment 0.1pct_PM

APPROVALS

| Signed by | Meaning of Signature | Server Date |
|------------------|--|-------------------------|
| Maureen Ramanan | Scanned Image is Identical to Original | 08-May-2018 12:49:24 PM |